

Yangians and quantizations of slices in the affine Grassmannian

Joel Kamnitzer

Dept. of Mathematics, University of Toronto; jkamnitz@math.toronto.edu

Ben Webster¹

Dept. of Mathematics, Northeastern University; b.webster@neu.edu

Alex Weekes

Dept. of Mathematics, University of Toronto; alex.weekes@utoronto.ca

Oded Yacobi

Dept. of Mathematics, University of Toronto; oyacobi@math.toronto.edu

Abstract. We study quantizations of transverse slices to Schubert varieties in the affine Grassmannian. The quantization is constructed using quantum groups called shifted Yangians — these are subalgebras of the Yangian we introduce which generalize the Brundan-Kleshchev shifted Yangian to arbitrary type. Building on ideas of Gerasimov-Kharchev-Lebedev-Oblezin, we prove that a quotient of the shifted Yangian quantizes a scheme supported on the transverse slices, and we formulate a conjectural description of the defining ideal of these slices which implies that the scheme is reduced. This conjecture also implies the conjectural quantization of the Zastava spaces for PGL_n of Finkelberg-Rybnykov.

1 Introduction

We initiate a program which relates the geometry of affine Grassmannians with the representation theory of shifted Yangians. More precisely, we study slices in affine Grassmannians which arise naturally in geometric representation theory; they correspond to weight spaces of irreducible representations under the geometric Satake correspondence. Our main result is that certain subquotients of Yangians quantize these slices.

There is a general program to study symplectic resolutions by means of the representation theory of their quantizations, generalizing the interplay between semisimple Lie algebras and nilpotent cones. We believe that the representation theory of shifted Yangians and its relationship to the geometry of slices in the affine Grassmannian will prove to be a very fruitful area of inquiry.

¹Supported by NSF grant DMS-1151473 and by NSA grant H98230-10-1-0199.

1.1 Slices in the affine Grassmannian

Let G be a complex semisimple group and consider its **thick affine Grassmannian** $\text{Gr} = G((t^{-1}))/G[t]$. Attached to each pair of dominant coweights $\lambda \geq \mu$, we have Schubert varieties $\text{Gr}^\lambda, \text{Gr}^\mu \subset \text{Gr}$, with $\text{Gr}^\mu \subset \overline{\text{Gr}^\lambda}$. The neighborhood in $\overline{\text{Gr}^\lambda}$ of a point in Gr^μ is encapsulated in a transversal slice to the latter variety in the former, which we denote by Gr_μ^λ . This slice is an important object of study in geometric representation theory because under the geometric Satake correspondence it is related to the μ weight space in the irreducible representation of G^\vee of highest weight λ .

The Manin triple $(\mathfrak{g}[t], t^{-1}\mathfrak{g}[[t^{-1}]], \mathfrak{g}((t^{-1})))$ provides Gr with the structure of a Poisson variety. The slice Gr_μ^λ is an affine Poisson subvariety and thus, its coordinate ring is naturally a Poisson algebra. The purpose of this paper is to explicitly describe quantizations of this Poisson algebra.

1.2 Quotients of shifted Yangians

The slice Gr_μ^λ is defined as the intersection $\overline{\text{Gr}^\lambda} \cap \text{Gr}_\mu$, where Gr_μ is an orbit of the group $G_1[[t^{-1}]]$, the first congruence subgroup of $G[[t^{-1}]]$. Thus on the level of functions $\mathcal{O}(\text{Gr}_\mu^\lambda)$ is a quotient of $\mathcal{O}(\text{Gr}^\mu)$, and $\mathcal{O}(\text{Gr}^\mu)$ is a subalgebra of $\mathcal{O}(G_1[[t^{-1}]])$. In order to quantize Gr_μ^λ we follow a three step procedure which mirrors this construction.

We first construct a version Y of the Yangian, which is a subalgebra of the Drinfeld Yangian. Next, we define natural subalgebras $Y_\mu \subset Y$, called **shifted Yangians**, quantize Gr_μ . This generalizes the shifted Yangian for \mathfrak{gl}_n introduced by Brundan-Kleshchev [BrKl1]. Finally, we define a quotient Y_μ^λ of Y_μ using some remarkable representations of Y as difference operators, constructed by Gerasimov-Kharchev-Lebedev-Oblezin [GKLO].

Theorem A. *The algebras defined above are all quantizations of the analogous geometric objects. That is:*

1. *The Yangian Y quantizes $G_1[[t^{-1}]]$.*
2. *The shifted Yangian Y_μ quantizes Gr_μ .*
3. *The quotient Y_μ^λ quantizes a (possibly non-reduced) scheme supported on Gr_μ^λ .*

Item (1) above is proven using the Drinfeld-Gavarini quantum groups duality, (2) follows simply from (1), and (3) follows using the GKLO representation. In fact, we produce a family $Y_\mu^\lambda(\mathbf{c})$ of quantizations which we conjecture to map surjectively to the universal family in the sense of Bezrukavnikov-Kaledin [BeKa].

Unfortunately, we are not able to prove that the scheme quantized by Y_μ^λ is reduced. However, we do provide a conjectural description of the generators of the ideal of Gr_μ^λ inside Gr_μ and prove that this conjecture implies that Y_μ^λ quantizes the reduced scheme structure on Gr_μ^λ . Moreover, we prove that this conjecture gives a simple description for the ideal defining Y_μ^λ .

1.3 Motivation and relation to other work

Brundan-Kleshchev [BrK11] construct an isomorphism between quotients of shifted Yangians of \mathfrak{gl}_n and W -algebras of \mathfrak{gl}_m . On one hand, it is known that W -algebras are quantizations of Slodowy slices. On the other hand, by the work of Mirković-Vybornov [MVy] we have an isomorphism between Slodowy slices for \mathfrak{sl}_m and slices in the affine Grassmannian for GL_n . Thus via these results, we see that quotients of shifted Yangians for \mathfrak{gl}_n quantize slices in the affine Grassmannian for GL_n . This motivated us to look for a direct construction of quantizations of affine Grassmannian slices (for any semisimple G) using quotients of shifted Yangians. (The idea that the Brundan-Kleshchev isomorphism should be thought as a quantization of the Mirković-Vybornov isomorphism was independently observed by Losev [Lo, Remark 5.3.4].)

If we take a limit of Gr_μ^λ as $\lambda \rightarrow \infty$ and $\lambda - \mu$ is fixed, then the slice Gr_μ^λ becomes the Zastava space $Z_{\lambda - \mu}$. Finkelberg-Rybnikov [FR] have given conjectural quantizations of Zastava spaces (for PGL_n) using quotients of Borel Yangians, which are a limit of shifted Yangians. Thus in this limit we prove their conjectures, dependent on the above mentioned conjecture about the ideal of Gr_μ^λ .

Earlier work on shifted Yangians by Brundan and Kleshchev [BrK12] suggest that one natural direction for future work is the study of a version of category \mathcal{O} over the algebra Y_μ^λ . Because of the geometric Satake correspondence, we think of category \mathcal{O} for Y_μ^λ as a categorification of a weight space in a representation of the Langlands dual group G^\vee . Thus we expect that these categories (with λ fixed) carry categorical \mathfrak{g}^\vee -actions. Moreover, conjectures of Braden, Licata, Proudfoot and the second author [BLPW] suggest that category \mathcal{O} for Y_μ^λ should be Koszul dual to similar categories constructed from quiver varieties (in type A, we expect that this reduces to parabolic-singular duality of Beilinson-Ginzburg-Soergel [BGS]).

1.4 Acknowledgements

We would like to thank Alexander Braverman, Pavel Etingof, Mikhail Finkelberg, Ivan Mirković, Sergey Oblezin, Travis Schedler and Catharina Stroppel for extremely useful conversations.

2 Symplectic structure on slices in the affine Grassmannian

2.1 Notation

For any group H , we will write $H((t^{-1})) = H(\mathbb{C}((t^{-1})))$ for its loop group and write $H[t] = H(\mathbb{C}[t])$ and $H[[t^{-1}]] = H(\mathbb{C}[[t^{-1}]])$ for its usual subgroups. Let $H_1[[t^{-1}]]$ denote the first congruence subgroup of $H[[t^{-1}]]$, i.e. the kernel of the evaluation at $t^{-1} = 0$, $H[[t^{-1}]] \rightarrow H$.

Throughout G will denote a fixed complex semisimple group, with opposite Borel subgroups B, B_- , unipotent subgroups N, N_- , maximal torus T , coweight lattice X , Weyl group W , set of roots Δ , simple roots $\{\alpha_i\}_{i \in I}$. We write $\{\omega_i\}_{i \in I}$ for the fundamental weights of the simply connected form of G .

Following Drinfeld, we use generators e_i, f_i, h_i for \mathfrak{g} where

$$[h_i, e_j] = (\alpha_i, \alpha_j) e_j, \quad [h_i, f_j] = -(\alpha_i, \alpha_j) f_j, \quad [e_i, f_j] = \delta_{ij} h_i$$

along with the usual Serre relations. Let $(a_{ij})_{1 \leq i, j \leq n}$ be the Cartan matrix of \mathfrak{g} , and let d_i be the unique coprime positive integers such that $b_{ij} = d_i a_{ij}$ is a symmetric matrix. Then the associated invariant form on \mathfrak{g} is defined by $(e_i, f_j) = \delta_{ij}$, and $(\alpha_i, \alpha_j) = (h_i, h_j) = d_i a_{ij}$, and in particular h_i is the image of α_i under the identification of \mathfrak{h} and \mathfrak{h}^* .

This is as opposed to the standard Chevalley generators e'_i, f'_i, h'_i , which we will identify as

$$e_i = -d_i^{1/2} e'_i, \quad f_i = -d_i^{1/2} f'_i, \quad h_i = d_i h'_i$$

In this way we have fundamental weights $\omega_i(h'_j) = \delta_{ij}$, and a lift of the Weyl group defined via $\overline{s}_i = \exp(f'_i) \exp(-e'_i) \exp(f'_i)$.

If μ is a weight or coweight, we write $\mu^* = -w_0 \mu$. Likewise, we write i^* if $\alpha_i^* = -w_0 \alpha_i$.

Let V be a representation of G , and let $v \in V, \beta \in W^*$. The matrix entry $\Delta_{\beta, v}$ is a function on G given by $\Delta_{\beta, v}(g) = \langle \beta, gv \rangle$. If $w_1, w_2 \in W$ and τ a dominant weight, we define

$$\Delta_{w_1 \tau, w_2 \tau}(g) = \langle \overline{w_1} v_{-\tau}, \overline{w_2} v_{\tau} \rangle$$

using the lift described above, where v_{τ} is the highest weight vector for the irreducible representation $V(\tau)$ and $v_{-\tau}$ is the dual lowest weight vector in $V(\tau^*)$.

Using this matrix entry (also known as generalized minor), we define the function $\Delta_{\beta, v}^{(s)}$ on $G((t^{-1}))$, for $s \in \mathbb{Z}$, whose value at g is the coefficient of the polynomial $\Delta_{\beta, v}(g)$. More precisely, these are given by the formula

$$\Delta_{\beta, v}(g) = \sum_{s=-\infty}^{\infty} \Delta_{\beta, v}^{(s)}(g) t^{-s}$$

2.2 Slices in the affine Grassmannian

Let G be a semisimple complex group. In this paper, we will work with the **thick affine Grassmannian** $\mathrm{Gr} = G((t^{-1}))/G[t]$. We have an embedding of the usual thin affine Grassmannian into the thick affine Grassmannian

$$G((t))/G[[t]] \cong G[t, t^{-1}]/G[t] \hookrightarrow G((t^{-1}))/G[t]$$

In this paper, we work with the thick affine Grassmannian since it is forced upon us by the non-commutative algebras we consider. One manifestation of this is the fact that the thick Grassmannian is an honest scheme, while the thin Grassmannian is only an ind-scheme. However, at a first reading, this difference will be of little importance, and the reader can pretend that we are working with the usual thin affine Grassmannian.

Any coweight λ can be thought of as $\mathbb{C}[t, t^{-1}]$ -point of G , which we can think of as a $\mathbb{C}((t^{-1}))$ -point as well. To avoid confusion, we use t^λ to denote this point in $G((t^{-1}))$. We also use t^λ for the image of t^λ in Gr .

Let λ and μ will denote dominant weights. Define

$$\mathrm{Gr}^\lambda = G[t]t^\lambda, \quad \mathrm{Gr}_\mu = G_1[[t^{-1}]]t^{w_0\mu}.$$

Recall that the thin affine Grassmannian is precisely $\cup_\lambda \mathrm{Gr}^\lambda$.

Our main object of interest will be

$$\mathrm{Gr}_\mu^\lambda := \overline{\mathrm{Gr}^\lambda} \cap \mathrm{Gr}_\mu.$$

This variety is a transverse slice to Gr^μ inside of $\overline{\mathrm{Gr}^\lambda}$ since Gr_μ intersects every Gr^ν transversely, and the intersection Gr_μ^μ is just the point $t^{w_0\mu}$. In particular, this variety is non-empty if and only if $\mu \leq \lambda$, that is, if $\mathrm{Gr}^\mu \subset \overline{\mathrm{Gr}^\lambda}$. These varieties arise naturally under the geometric Satake correspondence of Lusztig [Lu], Ginzburg [Gi], and Mirković-Vilonen [MV]: the intersection homology of Gr_μ^λ is identified with the μ weight space of the irreducible G^\vee -representation of highest weight λ .

Note that \mathbb{C}^\times acts on Gr by loop rotation. This action preserves the $G[t]$ and $G_1[[t^{-1}]]$ orbits and so \mathbb{C}^\times acts on Gr_μ^λ . The following result is standard.

Proposition 2.1.

1. Gr_μ^λ is an affine variety of dimension $2\langle \rho, \lambda - \mu \rangle$.
2. The action of \mathbb{C}^\times on Gr_μ^λ contracts Gr_μ^λ to the unique fixed point $t^{w_0\mu}$.

Example 2.2. If $\lambda = \mu + \alpha_i^\vee$, then Gr_μ^λ is isomorphic to the Kleinian singularity $\mathbb{C}^2/(\mathbb{Z}/n+2)$ where $n = \langle \mu, \alpha_i \rangle$. To see this, first we identify

$$\mathbb{C}^2/(\mathbb{Z}/n+2) = \{(u, v, w) : uv + w^{n+2} = 0\}$$

and then we define the isomorphism

$$\begin{aligned} \mathbb{C}^2/(\mathbb{Z}/n+2) &\rightarrow \text{Gr}_\mu^\lambda \\ (u, v, w) &\mapsto \phi_i \left(\begin{bmatrix} 1+wt^{-1}+\dots+w^{n+1}t^{-(n+1)} & vt^{-(n+1)} \\ ut^{-1} & 1-wt^{-1} \end{bmatrix} \right) t^{w_0\mu} \end{aligned}$$

where $\phi_i : SL_2 \rightarrow G$ denotes the SL_2 corresponding to α_i .

Let $G((t^{-1}))_\mu$ denote the stabilizer of $t^{w_0\mu}$ inside of $G((t^{-1}))$. The following easy result describes the stabilizer on the Lie algebra level.

Lemma 2.3. $\text{Lie}(G((t^{-1}))_\mu) = \mathfrak{t}[t] \oplus \bigoplus_{\alpha \in \Delta} t^{\langle \alpha, w_0\mu \rangle} \mathfrak{g}_\alpha[t]$.

Proof. The result follows immediately after observing that for $g \in G((t^{-1}))$, we have $g \in G((t^{-1}))_\mu$ if and only if $t^{-w_0\mu}gt^{w_0\mu} \in G[t]$. \square

In what follows, we will need the following set-theoretic description of $\overline{\text{Gr}}^\lambda$ due to Finkelberg-Mirković [FM]. As we shall see, it is much trickier to find a description of this variety with its natural reduced scheme structure.

Proposition 2.4. *Let $g \in G((t^{-1}))$. We have $[g] \in \overline{\text{Gr}}^\lambda$ if and only if $\Delta_{\beta,v}^{(s)}(g) = 0$ for all dominant weights τ , for all $v \in V(\tau)$, $\beta \in V(\tau)^*$ and for all $s < \langle \lambda, w_0\tau \rangle$.*

Proof. Fix τ and let k be the minimal s such that there exists $\beta \in V(\tau)^*$, $v \in V(\tau)$ with $\Delta_{\beta,v}^{(s)}(g) \neq 0$ (if such a minimum exists). It is easy to see that k only depends on the $G[t]$ double coset containing g . Thus if $[g] \in \overline{\text{Gr}}^\lambda$, we have that $k = \langle \lambda, w_0\tau \rangle$. The result follows. \square

The proof makes it clear that the Proposition holds even if τ only ranges over a set of dominant weights which spans (over \mathbb{Q}) the weight lattice.

2.3 Symplectic structure on the affine Grassmannian

There is a non-degenerate pairing on $\mathfrak{g}((t^{-1}))$ coming from residue and the invariant form on \mathfrak{g} . Hence the Lie algebras $\mathfrak{g}[t]$, $t^{-1}\mathfrak{g}[[t^{-1}]]$, and $\mathfrak{g}((t^{-1}))$ form a Manin triple (see [Dr1]). This induces a Poisson-Lie structure on $G((t^{-1}))$ with $G[t]$ and $G_1[[t^{-1}]]$ as Poisson subgroups. In particular, it coinduces a Poisson structure on Gr , by standard calculations which date back to work of Drinfeld [Dr2].

Let us state a couple of results concerning the interaction between this symplectic structure and the geometry considered in the previous section. These results were originally obtained by Mirković [Mir].

Theorem 2.5. *The subvarieties $\mathrm{Gr}_\mu^\lambda = \mathrm{Gr}^\lambda \cap \mathrm{Gr}_\mu$ are symplectic leaves of Gr .*

Proof. First we note that Gr_μ^λ are connected by [R, 1.4], since $\mathfrak{g}((t^{-1})) = \mathfrak{g}[t] \oplus t^{-1}\mathfrak{g}[[t^{-1}]]$. The argument is stated there for finite dimensional groups, but carries through to the loop situation without issues. Then the result follows from [LY, Corollary 2.9]. \square

These are not all symplectic leaves of Gr , since not every $G_1[[t^{-1}]]$ -orbit contains a point $t^{w_0\mu}$ and not every $G[t]$ orbit contains a point t^λ . A general symplectic leaf which lies in the thin affine Grassmannian is of the form $\mathrm{Gr}^\lambda \cap G_1[[t^{-1}]]gt^{w_0\mu}$ where $g \in G$.

Let $S^\mu = N((t^{-1}))t^\mu$. An **MV cycle** is a component of $\overline{\mathrm{Gr}^\lambda} \cap S^\mu$. By Mirković-Vilonen, these MV cycles give a basis for weight spaces of irreducible representations of the Langlands dual group. As we now see the MV cycles are Lagrangians in Gr_μ^λ .

Proposition 2.6. *$\overline{\mathrm{Gr}^\lambda} \cap S^\mu$ is a Lagrangian subvariety of Gr_μ^λ .*

Proof. First we prove that $\overline{\mathrm{Gr}^\lambda} \cap S^\mu \subset \mathrm{Gr}_\mu^\lambda$. Since N is unipotent, we have that $N((t^{-1})) = N_1[[t^{-1}]]N[t]$. Now by Lemma 2.3, we have that $N[t]t^{w_0\mu} = t^{w_0\mu}$. Hence $N((t^{-1}))t^\mu = N_1[[t^{-1}]]t^{w_0\mu}$ and thus $S^\mu \subset \mathrm{Gr}_\mu$.

From [MV], $\dim \overline{\mathrm{Gr}^\lambda} \cap S^\mu = \langle \rho, \lambda - \mu \rangle$ and thus the intersection $\overline{\mathrm{Gr}^\lambda} \cap S^\mu$ is half-dimensional in Gr_μ^λ . Hence it is Lagrangian if and only if it is coisotropic. The variety Gr_μ^λ is affine, and so it suffices to check that the Poisson bracket of any two functions that vanish on $\overline{\mathrm{Gr}^\lambda} \cap S^\mu$ vanishes there as well. The functions vanishing on $S^\mu \cap \overline{\mathrm{Gr}^\lambda}$ are generated by all functions of negative weight under the action of the coweight $\rho^\vee : \mathbb{C}^\times \rightarrow G$. Since that action preserves the Poisson structure, the Poisson bracket of two negative weight functions is again negative weight; this completes the proof. \square

It is natural to ask whether Gr_μ^λ has a symplectic resolution. Let us temporarily assume that G is of adjoint type and let us fix an sequence $\vec{\lambda} = (\lambda_1, \dots, \lambda_n)$ of fundamental coweights such that $\lambda = \lambda_1 + \dots + \lambda_n$. Then we have the open and closed convolutions

$$\mathrm{Gr}^{\vec{\lambda}} := \mathrm{Gr}^{\lambda_1} \tilde{\times} \dots \tilde{\times} \mathrm{Gr}^{\lambda_n}, \quad \overline{\mathrm{Gr}^{\vec{\lambda}}} := \overline{\mathrm{Gr}^{\lambda_1}} \tilde{\times} \dots \tilde{\times} \overline{\mathrm{Gr}^{\lambda_n}}$$

along with the convolution morphisms $m : \mathrm{Gr}^{\vec{\lambda}} \rightarrow \overline{\mathrm{Gr}^\lambda}$ and $\bar{m} : \overline{\mathrm{Gr}^{\vec{\lambda}}} \rightarrow \overline{\mathrm{Gr}^\lambda}$. Let

$$\mathrm{Gr}_\mu^{\vec{\lambda}} := m^{-1}(\mathrm{Gr}_\mu) \quad \overline{\mathrm{Gr}}_\mu^{\vec{\lambda}} := \bar{m}^{-1}(\mathrm{Gr}_\mu).$$

Recall that a normal variety X with a fixed symplectic structure Ω on its smooth locus is said to have **symplectic singularities** if, locally on X , there are resolutions of singularities $p: U \rightarrow X$ where $p^*\Omega$ is the restriction of a closed 2-form on U (which is not assumed to be non-degenerate on the exceptional locus).

A variety X is said to have **terminal singularities** if there is a resolution of singularities of X such that each irreducible exceptional fiber has positive discrepancy, that is, X is as close to being smoothly resolved as is crepantly possible. A **terminalization** $X \rightarrow Y$ is a map which is birational, proper, and crepant with X having terminal singularities. We say a variety X is **\mathbb{Q} -factorial** if every Weil divisor on X has an integer multiple which is Cartier.

Theorem 2.7. *The variety $\mathrm{Gr}_\mu^{\bar{\lambda}}$ has symplectic singularities, and $\mathrm{Gr}_\mu^{\bar{\bar{\lambda}}}$ is a \mathbb{Q} -factorial terminalization of $\mathrm{Gr}_\mu^{\bar{\lambda}}$.*

Proof. First, we claim that $\mathrm{Gr}_\mu^{\bar{\bar{\lambda}}}$ has singular locus in codimension ≥ 4 . Since Gr_μ is transverse to every $G[t]$ -orbit, the codimension of the singular locus cannot jump when we pass to $\mathrm{Gr}_\mu^{\bar{\bar{\lambda}}}$, so we need only establish the same result for $\mathrm{Gr}_\mu^{\bar{\lambda}}$, for which it suffices to consider the case of a fundamental coweight. If ω_i is a fundamental coweight, and ν is a dominant coweight such that $\mathrm{Gr}^\nu \subset \mathrm{Gr}^{\bar{\omega}_i}$, then we have that $\rho^\vee(\omega_i - \nu) \geq 2$, since $\omega_i - \alpha_j$ is never dominant. Thus, the singular locus Gr^ν has codimension at least 4.

As Beauville notes [Be, (1.2)], since $\mathrm{Gr}_\mu^{\bar{\bar{\lambda}}}$ is regular in codimension 3 and normal, the existence of a symplectic form on its smooth locus implies that it has symplectic singularities. Since we have a Poisson map $\mathrm{Gr}_\mu^{\bar{\bar{\lambda}}} \rightarrow \mathrm{Gr}_\mu^{\bar{\lambda}}$, this variety also has symplectic singularities. By a result of Namikawa [N1], this regularity in codimension 3 also implies that $\mathrm{Gr}_\mu^{\bar{\bar{\lambda}}}$ is terminal.

Since each local singularity in $\mathrm{Gr}_\mu^{\bar{\bar{\lambda}}}$ is a local singularity in $\mathrm{Gr}_\mu^{\bar{\lambda}}$, and these are the product of local singularities in $\mathrm{Gr}^{\bar{\omega}_i}$, we need only prove \mathbb{Q} -factoriality in this case. The group of Weil divisors of $\mathrm{Gr}^{\bar{\omega}_i}$ is the same as that of Gr^{ω_i} which is an affine bundle over G/P_i where P_i is the maximal parabolic containing all negative simple root spaces but $\mathfrak{g}_{-\alpha_i}$. Thus, the Weil divisor group of G/P_i is isomorphic to \mathbb{Z} .

Since $\mathrm{Gr}^{\bar{\omega}_i}$ is projective, *some* Weil divisor on $\mathrm{Gr}^{\bar{\omega}_i}$ is Cartier. Thus, the group generated by any non-trivial Weil divisor must intersect the image of the Cartier divisors, and so $\mathrm{Gr}^{\bar{\omega}_i}$ is \mathbb{Q} -factorial[†].

Furthermore, the map $\mathrm{Gr}_\mu^{\bar{\bar{\lambda}}} \rightarrow \mathrm{Gr}_\mu^{\bar{\lambda}}$ is well-known to be proper and birational. The preimage of Gr^μ for $\mu \neq \lambda, \lambda - \alpha_i$ has codimension ≥ 4 , so any exceptional divisor must be the closure of a component of the preimage of $\mathrm{Gr}^{\lambda - \alpha_i}$. The coefficients of

[†]We thank Alexander Braverman for suggesting this portion of the argument to us.

these divisors in the discrepancy can thus be computed locally in a neighborhood of $x \in \text{Gr}^{\lambda - \alpha_i}$, but the germ of the map is equivalent to the minimal resolution of a Kleinian singularity by Example 2.2. The Kleinian singularities are known to be crepant. \square

An obvious question is when $\text{Gr}_\mu^{\bar{\lambda}}$ has a symplectic resolution. First, we make the following conjecture.

Conjecture 2.8. *Any symplectic resolution of $\text{Gr}_\mu^{\bar{\lambda}}$ is of the form $\text{Gr}_\mu^{\bar{\bar{\lambda}}}$.*

We can easily see when $\text{Gr}_\mu^{\bar{\bar{\lambda}}}$ is actually a resolution.

Theorem 2.9. *The following are equivalent.*

1. $\text{Gr}_\mu^{\bar{\bar{\lambda}}}$ is smooth.
2. $\text{Gr}_\mu^{\bar{\lambda}} = \text{Gr}_\mu^{\bar{\bar{\lambda}}}$.
3. There do not exist coweights ν_1, \dots, ν_n such that $\nu_1 + \dots + \nu_n = \mu$, for all k , ν_k is a weight of $V(\lambda_k)$ and for some k , ν_k is not an extremal weight of $V(\lambda_k)$.

Proof. (i) \Rightarrow (ii): It is well-known that the smooth locus of $\overline{\text{Gr}^\lambda}$ is precisely Gr^λ . Thus the smooth locus of $\text{Gr}_\mu^{\bar{\bar{\lambda}}}$ is precisely $\text{Gr}_\mu^{\bar{\lambda}}$.

Now assume that there is a point x in $\text{Gr}_\mu^{\bar{\bar{\lambda}}}$ not in $\text{Gr}_\mu^{\bar{\lambda}}$; we know that $\text{Gr}_\mu^{\bar{\bar{\lambda}}}$ is not smooth at x . By the transversality of the $G_1[[t^{-1}]]$ and $G[t]$ orbits, the completion of $\text{Gr}_\mu^{\bar{\bar{\lambda}}}$ at x is the same as the completion of $\text{Gr}_\mu^{\bar{\lambda}}$ at x times something smooth. Therefore $\text{Gr}_\mu^{\bar{\bar{\lambda}}}$ cannot be smooth at x either.

(ii) \Rightarrow (i): clear.

(ii) \Rightarrow (iii): If there exist ν_1, \dots, ν_n as in (iii), then $(t^{\nu_1}, t^{\nu_1 + \nu_2}, \dots, t^\mu) \in \text{Gr}_\mu^{\bar{\bar{\lambda}}} \setminus \text{Gr}_\mu^{\bar{\lambda}}$.

(iii) \Rightarrow (ii): Suppose that there exists

$$(L_1, \dots, L_n) \in \text{Gr}_\mu^{\bar{\bar{\lambda}}} \setminus \text{Gr}_\mu^{\bar{\lambda}}$$

Recall that we have a $\mathbb{C}^\times \times T$ action on Gr where the first factor acts by loop rotation. Consider a map $\mathbb{C}^\times \rightarrow \mathbb{C}^\times \times T$ which is the identity into the first factor and a generic dominant coweight into the second factor. We get a resulting \mathbb{C}^\times action on Gr whose attracting sets are the I_- orbits, where I_- is the preimage of B under $G[[t^{-1}]] \rightarrow G$.

Let $(t^{\mu_1}, \dots, t^{\mu_n}) = \lim_{s \rightarrow 0} s \cdot (L_1, \dots, L_n)$. From the definition of $\text{Gr}_\mu^{\bar{\bar{\lambda}}}$, we see that $\mu_n = \mu$. Also, we see that for each k , $d(t^{\mu_{k-1}}, t^{\mu_k}) \leq \lambda_k$ (where $d : \text{Gr} \times \text{Gr} \rightarrow$

X_+ denotes the $G((t^{-1}))$ -invariant distance function on Gr and so $\nu_k := \mu_k - \mu_{k-1}$ is a weight of $V(\lambda_k)$. Thus we obtain ν_1, \dots, ν_n with $\nu_1 + \dots + \nu_n = \mu$. Moreover, since $(L_1, \dots, L_n) \notin \text{Gr}_\mu^{\bar{\lambda}}$, for some k , $d(L_{k-1}, L_k) < \lambda_k$ and so ν_k is a non-extremal weight of $V(\lambda_k)$. \square

If λ is a sum of minuscule coweights, then the above conditions hold. For any simple G not of type A, there are non-minuscule fundamental coweights λ ; for such λ , we can choose μ such that the above conditions do not hold. So we expect that there exist $\text{Gr}_\mu^{\bar{\lambda}}$ which do not admit symplectic resolutions (and which we know do not admit ones of the form $\text{Gr}_\mu^{\bar{\lambda}}$).

2.4 Beilinson-Drinfeld Grassmannian

Using the Beilinson-Drinfeld Grassmannian, we can define a family of Poisson varieties over \mathbb{A}^n whose special fibre is $\text{Gr}_\mu^{\bar{\lambda}}$. In this work, this family will only be used as motivation for a similar family of quantizations of $\text{Gr}_\mu^{\bar{\lambda}}$; as illustrated in works such as [BeKa, BPW, Lo], the universal symplectic deformation of a symplectic singularity as a symplectic variety is intimately tied to understanding its quantizations (see section 4.4). From this perspective, a natural next step (beyond the scope of this paper) would be to study quantizations of the total spaces of these deformations, not just of a single fibre.

Recall that we have the moduli interpretation of the affine Grassmannian

$$\text{Gr} = \{(E, \phi) : E \text{ is a principal } G\text{-bundle on } \mathbb{P}^1 \text{ and} \\ \phi : E|_{\mathbb{P}^1 \setminus \{0\}} \rightarrow E^0|_{\mathbb{P}^1 \setminus \{0\}} \text{ is an isomorphism}\}$$

where E^0 denotes the trivial G -bundle. We say that (E, ϕ) has **Hecke type** λ at 0 if (E, ϕ) gives a point in Gr^λ under the above identification.

Note that the action of $G[[t^{-1}]]$ by left multiplication in the homogeneous space definition becomes change of trivialization in the new definition. Thus the $G[[t^{-1}]]$ orbit of (E, ϕ) is determined by isomorphism class of the G -bundle E , which is given by a dominant coweight. Note also that the action of $G_1[[t^{-1}]]$ corresponds to changes of trivialization which do not change anything at ∞ .

Let μ be a dominant weight and let P be the corresponding standard parabolic subgroup (so that W_P is the stabilizer of μ in the Weyl group). Let E be a principal G -bundle of type μ . Then E has a canonical P -structure.

Now let $(E, \phi) \in \text{Gr}$. Let μ be the isomorphism type of E . Then ϕ_∞ carries the parabolic structure at ∞ to a parabolic subgroup of G of type μ . Hence we see that the $G_1[[t^{-1}]]$ orbits on Gr are labelled by a pair consisting of a dominant weight μ and a parabolic subgroup of G of type μ . In particular Gr_μ is the locus of those (E, ϕ) where E has isomorphism type μ and the parabolic subgroup produced is the standard one.

We now will consider the Beilinson-Drinfeld deformation of the affine Grassmannian. This is a family $\text{Gr}_{\mathbb{A}^n}$ over \mathbb{A}^n whose fibre at $a_1, \dots, a_n \in \mathbb{A}^n$ is given as follows:

$$\text{Gr}_{a_1, \dots, a_n} = \{(E, \phi) : E \text{ is a principal } G\text{-bundle on } \mathbb{P}^1 \text{ and} \\ \phi : E|_{\mathbb{P}^1 \setminus \{a_1, \dots, a_n\}} \rightarrow E^0|_{\mathbb{P}^1 \setminus \{a_1, \dots, a_n\}} \text{ is an isomorphism} \}$$

Let $\text{Gr}_{\mu, \mathbb{A}^n}$ be the locus of (E, ϕ) where E has isomorphism type μ and the parabolic subgroup at ∞ is the standard one.

Specializing to one choice of parameters, we can consider changes of trivialization acting on $\text{Gr}_{a_1, \dots, a_n}$. Let $G_1(\mathbb{P}^1 \setminus \{a_1, \dots, a_n\})$ denote the kernel of $G(\mathbb{P}^1 \setminus \{a_1, \dots, a_n\}) \rightarrow G$ given by evaluation at ∞ . Then, $\text{Gr}_{\mu, (a_1, \dots, a_n)}$ is an orbit of $G_1(\mathbb{P}^1 \setminus \{a_1, \dots, a_n\})$.

We may also think of this locus in terms of the \mathbb{C}^\times action. We have an action of \mathbb{C}^\times on $\text{Gr}_{\mathbb{A}^n}$ coming from the action of \mathbb{C}^\times on \mathbb{P}^1 . Note that this action moves the base \mathbb{A}^n . On the central fibre $\text{Gr}_{(0, \dots, 0)} = \text{Gr}$ this action of \mathbb{C}^\times restricts to the loop rotation action on Gr . Hence the fixed points of this \mathbb{C}^\times action are the same as the fixed points of the loop rotation action, namely the sets Gt^μ inside the affine Grassmannian. Moreover, we have that $\text{Gr}_{\mu, \mathbb{A}^n}$ is the attracting set for $t^{w_0\mu}$ under the \mathbb{C}^\times action.

We have a fiberwise Poisson structure on $\text{Gr}_{\mathbb{A}^n}$ using the Manin triples described in Etingof-Kazhdan [EK], Corollary 2.10 and Proposition 2.12. As in Section 2.3, we get a Poisson structure on $\text{Gr}_{\mu, (a_1, \dots, a_n)}$.

Now, let us choose an expression $\lambda = \lambda_1 + \dots + \lambda_n$, where $\lambda_1, \dots, \lambda_n$ are fundamental coweights. This gives us an X_+ colored divisor D on \mathbb{P}^1 defined by $D = \sum \lambda_i a_i$. We will think of D as a function $\mathbb{P}^1 \rightarrow X_+$. Now we define

$$\text{Gr}_{\mu, (a_1, \dots, a_n)}^{\lambda_1, \dots, \lambda_n} := \{(E, \phi) \in \text{Gr}_{\mu, (a_1, \dots, a_n)} : (E, \phi) \text{ has Hecke type } D(x) \text{ for all } x \in \mathbb{P}^1\}$$

From the above analysis, it is possible to show that these are symplectic leaves in $\text{Gr}_{\mu, (a_1, \dots, a_n)}$.

Fixing $(\lambda_1, \dots, \lambda_n)$ and letting (a_1, \dots, a_n) vary, this forms a family of \mathbb{A}^n . The central fibre of this family is Gr_μ^λ .

Now, define

$$\overline{\text{Gr}_{\mu, (a_1, \dots, a_n)}^{\lambda_1, \dots, \lambda_n}} := \{(E, \phi) \in \text{Gr}_{\mu, (a_1, \dots, a_n)} : (E, \phi) \text{ has Hecke type } \leq D(x) \text{ for all } x \in \mathbb{P}^1\}$$

Then we obtain a flat family of symplectic varieties over \mathbb{A}^n whose central fibre is $\text{Gr}_\mu^{\bar{\lambda}}$.

2.5 Direct system on slices and Zastava spaces

We will now look at what happens to $\text{Gr}_\mu^{\bar{\lambda}}$ when we increase λ, μ , keeping $\lambda - \mu$ fixed.

Let us fix ν in the positive coroot cone. Let μ, μ' be dominant coweights with $\mu' - \mu$ dominant. From Lemma 2.3, the stabilizer of $t^{w_0\mu'}$ in $G_1[[t^{-1}]]$ contains the stabilizer of $t^{w_0\mu}$ in $G_1[[t^{-1}]]$. So we can define a map $\text{Gr}_\mu \rightarrow \text{Gr}_{\mu'}$ by $gt^{w_0\mu} \mapsto gt^{w_0\mu'}$. From Proposition 2.4, we see that this restricts to a map $\overline{\text{Gr}}_\mu^{\mu+\nu} \rightarrow \overline{\text{Gr}}_{\mu'}^{\mu'+\nu}$. By construction, it is a Poisson map.

Clearly these maps are compatible with composition. Thus with ν fixed we get a direct system of slices $\left\{ \overline{\text{Gr}}_\mu^{\mu+\nu} \right\}_\mu$. The limit of this system is an ind-scheme, but in general it will not be represented by a scheme.

On the other hand, we can consider the Zastava space Z_ν , an affine variety, as defined in [FM]. It is a compactification of the moduli space Z_ν° of based maps from \mathbb{P}^1 into G/B of degree ν . The variety Z_ν carries an action of \mathbb{C}^\times , extending the action of \mathbb{C}^\times on Z_ν° which rotates the source of the map.

The following result is Theorem 2.8 from Braverman-Finkelberg [BF]. It shows that the algebras of functions $\mathcal{O}(\overline{\text{Gr}}_\mu^{\mu+\nu})$ stabilize to $\mathcal{O}(Z_\nu)$.

Theorem 2.10. *There exists a map $\overline{\text{Gr}}_\mu^{\mu+\nu} \rightarrow Z_\nu$. These maps are compatible with the above direct system on the slices and with the actions of \mathbb{C}^\times . Moreover, the induced maps $\mathcal{O}(Z_\nu)_N \rightarrow \mathcal{O}(\overline{\text{Gr}}_\mu^{\mu+\nu})_N$ are isomorphisms if $N \leq \langle \alpha_i, \mu \rangle$ for all i .*

Remark 2.11. The theorem provides Z_ν with a Poisson structure. On the other hand, Z_ν° carries a symplectic structure as described in [FKMM]. It is expected that these two structures are compatible.

Example 2.12. Let us take $G = PGL_2$ and $\nu = \alpha^\vee$, the simple coroot. Then (as in Example 2.2), for $n \geq 0$,

$$\overline{\text{Gr}}_{n\omega^\vee}^{n\omega^\vee + \alpha^\vee} \cong \{(u, v, w) : uv + w^{n+2} = 0\}$$

Moreover, for $m \geq n$, the map $\overline{\text{Gr}}_{n\omega^\vee}^{n\omega^\vee + \alpha^\vee} \rightarrow \overline{\text{Gr}}_{m\omega^\vee}^{m\omega^\vee + \alpha^\vee}$ is given by $(u, v, w) \mapsto (u, vw^{m-n}, w)$. This is because we have an equality in Gr_{PGL_2}

$$\begin{bmatrix} 1+wt^{-1}+\dots+w^{n-1}t^{-(n+1)} & vt^{-(n+1)} \\ ut^{-1} & 1-wt^{-1} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & t^m \end{bmatrix} = \begin{bmatrix} 1+wt^{-1}+\dots+w^{m+1}t^{-(m+1)} & vw^{m-n}t^{-(m+1)} \\ ut^{-1} & 1-wt^{-1} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & t^m \end{bmatrix}.$$

On the other hand, the Zastava space Z_α is \mathbb{A}^2 . The map in Theorem 2.10 is given by $(u, v, w) \mapsto (u, w)$.

With respect to the \mathbb{C}^\times action on

$$\overline{\text{Gr}}_{n\omega^\vee}^{n\omega^\vee + \alpha^\vee} = \{(u, v, w) : uv + w^{n+2} = 0\}$$

the variables u, w have weight 1 and v has weight $n+1$. So, we can see that

$$\mathcal{O}(Z_\alpha) = \mathbb{C}[u, w] \rightarrow \mathcal{O}(\overline{\text{Gr}}_{n\omega^\vee}^{n\omega^\vee + \alpha^\vee}) = \mathbb{C}[u, v, w]/(uv + w^{n+2})$$

is an isomorphism in degrees $0, \dots, n$ as predicted by Theorem 2.10.

The Poisson structure on $\text{Gr}_{n\omega^\vee}^{\overline{n\omega^\vee + \alpha^\vee}}$ is given by

$$\{w, u\} = u \quad \{w, v\} = -v \quad \{u, v\} = (n+2)w^{n+1}$$

while the Poisson structure on Z_α is given by

$$\{w, u\} = u.$$

Finally, note that the \mathbb{C} -points of the ind-scheme $\lim_n \text{Gr}_{n\omega^\vee}^{\overline{n\omega^\vee + \alpha^\vee}}$ are

$$\{(u, w) : u \in \mathbb{C}^\times, w \in \mathbb{C}\} \cup \{(0, 0)\}$$

which is a proper subset of \mathbb{C}^2 and hence this ind-scheme is not equal to \mathbb{A}^2 .

2.6 Description of the Poisson structure

We would like to describe the Poisson structure on $G_1[[t^{-1}]]$ in a little more detail. Let $C \in \mathfrak{g} \otimes \mathfrak{g}$ be the Casimir element for the bilinear form. Picking dual bases, we may represent this element as $C = \sum J_a \otimes J^a$; this Casimir element allows us to describe the Poisson bracket of two minors. This can be written more compactly using the series $\Delta_{\beta, v}(u) = \sum_{s \geq 0} \Delta_{\beta, v}^{(s)} u^{-s}$. Note that $\Delta_{\beta, v}^{(0)} = \langle \beta, w \rangle$ is a constant function.

Proposition 2.13. *In $\mathcal{O}(G_1[[t^{-1}]])[[u_1^{-1}, u_2^{-1}]]$, the Poisson bracket $\{\Delta_{\beta_1, v_1}(u_1), \Delta_{\beta_2, v_2}(u_2)\}$ is equal to*

$$\frac{1}{u_1 - u_2} \sum_a \Delta_{\beta_1, J_a v_1}(u_1) \Delta_{\beta_2, J^a v_2}(u_2) - \Delta_{J_a \beta_1, v_1}(u_1) \Delta_{J^a \beta_2, v_2}(u_2)$$

Proof. The cobracket $\mathfrak{g}((t^{-1})) \rightarrow \mathfrak{g}((u_1)) \otimes \mathfrak{g}((u_2))$ is coboundary. If we let $r(u_1, u_2) = \frac{C}{u_1 - u_2}$ it is given by

$$a(t) \mapsto [a(u_1) \otimes 1 + 1 \otimes a(u_2), r(u_1, u_2)].$$

As described earlier, the Lie algebra $\mathfrak{g}((t^{-1}))$ carries an inner product $(f, g)_t = -\text{res}_{t=0}(f, g)$ for which $t^{-1}\mathfrak{g}[[t^{-1}]]$ is Lagrangian and complementary to $\mathfrak{g}[t]$; this realizes $\mathfrak{g}((t^{-1}))$ as the (topological) Drinfeld double of $t^{-1}\mathfrak{g}[[t^{-1}]]$. In particular, $G_1[[t^{-1}]] \subset G((t^{-1}))$ is a Poisson subgroup, and the Poisson bracket of any two functions on $G_1[[t^{-1}]]$ can be calculated taking the bracket of any two extensions to all of $G((t^{-1}))$ and then restricting to $G_1[[t^{-1}]]$.

Thus, the Poisson structure on $G((t^{-1}))$ is defined by $r^L(u_1, u_2) - r^R(u_1, u_2)$, the difference of the left translation and right translation of the element $r(u_1, u_2)$

considered as a bivector at the identity. If $X \in t^{-1}\mathfrak{g}[[t^{-1}]]$, and $g \in G_1[[t^{-1}]]$, we identify X with a tangent vector at g by left translation. Then we have

$$(d\Delta_{\beta,v})_g(X) = \langle \beta, gXv \rangle.$$

Hence

$$\begin{aligned} & \{\Delta_{\beta_1,v_1}(u_1), \Delta_{\beta_2,v_2}(u_2)\}(g) = (d\Delta_{\beta_1,v_1})_g \otimes (d\Delta_{\beta_2,v_2})_g(g) \\ &= \frac{1}{u_1 - u_2} \left(\sum_a \langle \beta_1, g(u_1)J_a v_1 \rangle \langle \beta_2, g(u_2)J^a v_2 \rangle - \sum_a \langle \beta_1, J_a g(u_1)v_1 \rangle \langle \beta_2, J^a g(u_2)v_2 \rangle \right) \end{aligned}$$

and then the proposition follows from the invariant of the pairing between dual representations. \square

We can unpack Proposition 2.13 into the following equations:

$$\{\Delta_{\beta_1,v_1}^{(r+1)}, \Delta_{\beta_2,v_2}^{(s)}\} - \{\Delta_{\beta_1,v_1}^{(r)}, \Delta_{\beta_2,v_2}^{(s+1)}\} = \sum \Delta_{J_a \beta_1, v_1}^{(r)} \Delta_{J^a \beta_2, v_2}^{(s)} - \Delta_{\beta_1, J_a v_1}^{(r)} \Delta_{\beta_2, J^a v_2}^{(s)} \quad (1)$$

for $r, s \geq 0$. These equations specify all the desired Poisson brackets.

2.7 A conjectural description of the ideal of Gr_μ^λ

In this section, we describe a conjectural description of the ideal of Gr_μ^λ as a subvariety of $\text{Gr}_0 = G_1[[t^{-1}]]$. Let G_{sc} denote the simply connected cover of G . Note that the natural map $G_{sc}[[t^{-1}]]_1 \rightarrow G[[t^{-1}]]_1$ is an isomorphism. This allows us to consider $\Delta_{\omega_i, \omega_i}^{(s)}$ as functions on $G_1[[t^{-1}]]$, even if ω_i are not weights of G (for example if G is of adjoint type).

We begin with the case of $\mu = 0$. Let J_0^λ denote the ideal in $\mathcal{O}(G_1[[t^{-1}]])$ Poisson generated by $\Delta_{\omega_i, \omega_i}^{(s)}$ for $s > \langle \lambda, \omega_{i^*} \rangle$ and for $i \in I$.

Conjecture 2.14. *The ideal of Gr_0^λ in $\mathcal{O}(G_1[[t^{-1}]])$ is J_0^λ .*

Let us make some comments on this conjecture. First, we have the following result.

Proposition 2.15. *J_0^λ is generated as an ordinary ideal by $\Delta_{\beta,v}^{(s)}$ for $s > \langle \lambda, \omega_{i^*} \rangle$ and for $i \in I$ and where β, v range over bases for $V(\omega_i)^*$ and $V(\omega_i)$.*

Proof. Let I be the ideal generated as an ordinary ideal by $\Delta_{\beta,v}^{(s)}$ for $s > \langle \lambda, \omega_{i^*} \rangle$. First, we show that this ideal is contained in J_0^λ .

First, we claim that $\Delta_{\omega_i, v}^{(s)} \in J_0^\lambda$ for all $v \in V(\omega_i)$ and $s > \langle \lambda, \omega_{i^*} \rangle$. We proceed by downward induction on the weight of v . The base case of v is highest weight

follows by definition. For the inductive step, suppose that v is not highest weight. In this case, $v = \sum f_j v_j$ for some v_j of higher weight than v .

Fix s with $s > \langle \lambda, \omega_{i^*} \rangle$. Using (1) and the expression for the Casimir (for notation see Section 3.2)

$$C = C_{\mathfrak{h}} + \sum_{\alpha \in \Phi_+} C_{\alpha} e_{\alpha} \otimes f_{\alpha} + C_{\alpha} f_{\alpha} \otimes e_{\alpha},$$

where $(e_{\alpha}, f_{\alpha}) = C_{\alpha}^{-1}$, we see that

$$\{\Delta_{\omega_i, v_j}^{(s)}, \Delta_{\omega_j, s_j \omega_j}^{(1)}\} = -\Delta_{\beta, f_j v_j}^{(s)}$$

Thus we see that

$$\Delta_{\omega_i, v}^{(s)} = \sum_j \Delta_{\omega_i, f_j v_j}^{(s)} = - \sum_j \{\Delta_{\omega_i, v_j}^{(s)}, \Delta_{\omega_j, s_j \omega_j}^{(1)}\}.$$

All the terms on the right hand side lie in J_0^{λ} by the inductive assumption, and thus $\Delta_{\omega_i, v}^{(s)} \in J_0^{\lambda}$.

Now we claim that $\Delta_{\beta, v}^{(s)} \in J_0^{\lambda}$ for all $\beta \in V(\omega_i)^*$, $v \in V(\omega_i)$ and $s > \langle \lambda, \omega_{i^*} \rangle$.

We have already proven this claim when $\beta = v_{-\omega_i}$, so we proceed by induction on the weight of β . Suppose that $\beta \in V(\omega_i)^*$ is not lowest weight and assume that the claim holds for all β of lower weight. In this case, we can write $\beta = \sum e_j \beta_j$ for some β_j of lower weight.

Fix s with $s > \langle \lambda, \omega_{i^*} \rangle$. Again using the above expression for the Casimir we find that

$$\{\Delta_{\beta_j, v}^{(s)}, \Delta_{s_j \omega_j, \omega_j}^{(1)}\} = \Delta_{\beta_j, e_j v}^{(s)} - \Delta_{e_j \beta_j, v}^{(s)},$$

Thus we see that

$$\Delta_{\beta, v}^{(s)} = \sum_j \Delta_{e_j \beta_j, v}^{(s)} = \sum_j \{\Delta_{\beta_j, v}^{(s)}, \Delta_{s_j \omega_j, \omega_j}^{(1)}\} - \Delta_{\beta_j, e_j v}^{(s)}.$$

All the terms on the right hand side lie in J_0^{λ} by the inductive assumption, and thus $\Delta_{\beta, v}^{(s)} \in J_0^{\lambda}$. This shows that $I \subset J_{\lambda}^0$.

It remains to show that I is a Poisson ideal. Since $\Delta_{\beta, v}^{(s)}$, for $\beta \in V(\omega_i)^*$, $v \in V(\omega_i)$, $i \in I$, generates $\mathcal{O}(G_1[[t^{-1}]])$, it suffices to check that I is closed under Poisson bracket with these elements. This follows immediately from (1). \square

Combining this proposition with Proposition 2.4, we obtain the following.

Corollary 2.16. *The vanishing set of J_0^{λ} is $\text{Gr}_0^{\bar{\lambda}}$.*

Thus in order to establish Conjecture 2.14, it only remains to show that I_0^{λ} is radical.

Remark 2.17. Let $G = SL_n$. By an observation which goes back to Lusztig, we know that there is an isomorphism $\text{Gr}_0^{\overline{n\omega_1}} \cong \mathcal{N}$, the nilpotent cone of \mathfrak{sl}_n . For any dominant coweight λ with $\lambda \leq n\omega_1$, under this isomorphism $\text{Gr}_0^{\overline{\lambda}}$ is taken to a nilpotent orbit closure. Thus, the above conjecture implies generators for the ideal of a nilpotent orbit closure inside the nilpotent cone of \mathfrak{sl}_n . From this perspective, one can see that Conjecture 2.14 would imply the main result of Weyman [W], which gives generators for the ideals of nilpotent orbit closures. This gives additional evidence toward the conjecture, but also suggests it will be difficult to prove.

Remark 2.18. One could imagine a similar conjecture for the ideal of $\text{Gr}^{\overline{\lambda}}$ inside of the homogeneous coordinate ring of Gr . However, this conjecture is false, already for SL_2 and $\lambda = \alpha$.

We will need the following generalization of Conjecture 2.14 which describes the ideal of $\text{Gr}_\mu^{\overline{\lambda}}$. Consider the subgroup $G_1[[t^{-1}]]_\mu$ defined as the stabilizer in $G_1[[t^{-1}]]$ of $t^{w_0\mu}$. Note that Lemma 2.3, $G_1[[t^{-1}]]_\mu \subset N_1[[t^{-1}]]$.

By the orbit-stabilizer theorem, we see that $\text{Gr}_\mu = G_1[[t^{-1}]]/G_1[[t^{-1}]]_\mu$ and so $\mathcal{O}(\text{Gr}_\mu) = \mathcal{O}(G_1[[t^{-1}]]^{G_1[[t^{-1}]]_\mu})$. Moreover the map $G_1[[t^{-1}]] \rightarrow \text{Gr}_\mu$ is Poisson and thus $\mathcal{O}(\text{Gr}_\mu)$ is a Poisson subalgebra of $\mathcal{O}(G_1[[t^{-1}]])$.

Lemma 2.19. *The subalgebra $\mathcal{O}(\text{Gr}_\mu)$ contains*

$$\begin{aligned} &\Delta_{s_i\omega_i, \omega_i}^{(s)}, \text{ for all } i \in I, s > 0, \quad \Delta_{\omega_i, \omega_i}^{(s)}, \text{ for all } i \in I, s > 0, \\ &(\Delta_{\omega_i, s_i\omega_i} / \Delta_{\omega_i, \omega_i})^{(s)}, \text{ for all } i \in I, s > \langle \mu^*, \alpha_i \rangle \end{aligned}$$

Later we will see that these elements generate $\mathcal{O}(\text{Gr}_\mu)$ as a Poisson algebra.

Proof. Note that the action of $G_1[[t^{-1}]]_\mu$ on $\mathcal{O}(G_1[[t^{-1}]])$ is given by $(k \cdot f)(g) = g(fk)$ for $k \in G_1[[t^{-1}]]_\mu$, $f \in \mathcal{O}(G_1[[t^{-1}]])$ and $g \in G_1[[t^{-1}]]$. In particular, we see that $k \cdot \Delta_{\beta, v} = \Delta_{\beta, kv}$.

Since $G_1[[t^{-1}]]_\mu \subset N_1[[t^{-1}]]$, the minors $\Delta_{\omega_i, \omega_i}$ and $\Delta_{s_i\omega_i, \omega_i}$ will be $G_1[[t^{-1}]]_\mu$ -invariant. Hence all $\Delta_{s_i\omega_i, \omega_i}^{(s)}$, $\Delta_{\omega_i, \omega_i}^{(s)}$ all lie in Gr_μ .

On the other hand, let us consider the coefficients of the $\Delta_{\omega_i, s_i\omega_i}$ minor. If $k \in G_1[[t^{-1}]]_\mu$, then we have $k \cdot v_{s_i\omega_i} = v_{s_i\omega_i} + \Delta_{\omega_i, s_i\omega_i}(k)v_{\omega_i}$. Hence if $g \in G_1[[t^{-1}]]$, then

$$\frac{\Delta_{\omega_i, s_i\omega_i}(gk)}{\Delta_{\omega_i, \omega_i}(gk)} = \frac{\Delta_{\omega_i, s_i\omega_i}(g) + \Delta_{\omega_i, \omega_i}(g)\Delta_{\omega_i, s_i\omega_i}(k)}{\Delta_{\omega_i, \omega_i}(g)} = \frac{\Delta_{\omega_i, s_i\omega_i}(g)}{\Delta_{\omega_i, \omega_i}(g)} + \Delta_{\omega_i, s_i\omega_i}(k)$$

By Lemma 2.3, we have $\text{val } \Delta_{\omega_i, s_i\omega_i}(k) \geq \langle w_0\mu, \alpha_i \rangle$. Hence the coefficient of t^{-s} in $\Delta_{\omega_i, s_i\omega_i} / \Delta_{\omega_i, \omega_i}$ is invariant under the action of $G_1[[t^{-1}]]_\mu$ for $s > \langle \mu^*, \alpha_i \rangle$. Thus $(\Delta_{\omega_i, s_i\omega_i} / \Delta_{\omega_i, \omega_i})^{(s)} \in \mathcal{O}(\text{Gr}_\mu)$ for $s > \langle \mu^*, \alpha_i \rangle$. \square

Let J_μ^λ denote the ideal of $\mathcal{O}(\text{Gr}_\mu)$ Poisson generated by $\Delta_{\omega_i, \omega_i}^{(s)}$ for $i \in I$ and $s > \langle \lambda - \mu, \omega_i^* \rangle = m_i$.

Conjecture 2.20. *The ideal of Gr_μ^λ in $\mathcal{O}(\text{Gr}_\mu)$ is J_μ^λ .*

This conjecture generalizes Conjecture 2.14. When $\mu \neq 0$, we do not have a set of (ordinary) generators for J_μ^λ as in Proposition 2.15. However, we will now establish an analog of Corollary 2.16.

Proposition 2.21. *The vanishing locus of J_μ^λ is Gr_μ^λ .*

Proof. The vanishing locus of J_μ^λ is the union of the symplectic leaves in the vanishing locus of $\Delta_{\omega_i, \omega_i}^{(s)}$ for $i \in I$ and $s > \langle \lambda - \mu, \omega_i^* \rangle = m_i$; after all, the vanishing set is a union of symplectic leaves and if these functions vanish on a symplectic leaf, so do all Poisson brackets with them.

These generalized minors vanish on Gr_μ^λ by Proposition 2.4. So it suffices to prove the vanishing locus of our generators does not contain Gr_μ^ν for some $\nu \not\leq \lambda$.

Fix $\nu \not\leq \lambda$ such that $\mu \leq \nu$. Then for some i , then $d = \langle \nu - \mu, \omega_i^* \rangle > \langle \lambda - \mu, \omega_i^* \rangle$. We will prove that there exists a point in Gr_μ^ν on which $\Delta_{\omega_i, \omega_i}^{(d)}$ is non-zero.

Let $I_+^+ = I \subset G((t^{-1}))$ denote the standard Iwahori and let $I_-^+ = w_0 I_+^+ w_0^{-1}$ be the preimage of B_- in $G[t]$. We claim that it suffices to prove that

$$I_-^+ t^{w_0 \nu} I_+^+ \cap G_-[[t^{-1}]] t^{w_0 \mu} \neq \emptyset \text{ in } G((t^{-1})) \quad (2)$$

To see that (2) suffices, let $g \in G_1[[t^{-1}]]$ such that $gt^{w_0 \mu}$ lies in the above intersection. As, $I_-^+, I_+^+ \subset G[t]$, we see that $gt^{w_0 \mu} \in \text{Gr}_\mu^\nu$. Finally, we can write $g = b_- t^{w_0 \nu} b_+ t^{-w_0 \mu}$ for $b_- \in I_-^+, b_+ \in I_+^+$ and an elementary computation shows that $\Delta_{\omega_i, \omega_i}^{(d)}(b_- t^{w_0 \nu} b_+ t^{-w_0 \mu}) \neq 0$.

To prove (2), we work in the affine flag variety $G((t^{-1}))/I$ and note that (2) is equivalent to non-emptiness of the intersection $I_-^+ t^{w_0 \nu} \cap G_-[[t^{-1}]] t^{w_0 \mu}$ in $G((t^{-1}))/I$. Let I_+^- denote the preimage of B in $G[[t^{-1}]]$ under evaluation at $t^{-1} = 0$. Since μ is dominant, B fixes $t^{w_0 \mu}$ and thus $G_-[[t^{-1}]] t^{w_0 \mu} = I_+^- t^{w_0 \mu}$. Thus we reduce to proving that

$$I_-^+ t^{w_0 \nu} \cap I_+^- t^{w_0 \mu} \neq \emptyset \text{ in } G((t^{-1}))/I.$$

Twisting by w_0 , we reduce to proving that

$$I_+^+ w_0 t^{w_0 \nu} \cap I_-^- w_0 t^{w_0 \mu} \neq \emptyset \text{ in } G((t^{-1}))/I.$$

where I_-^- is the preimage of B_- in $G[[t^{-1}]]$. From general theory of flag varieties, this is equivalent to $w_0 t^{w_0 \nu} \geq w_0 t^{w_0 \mu}$ in the Bruhat order on the (extended) affine Weyl group. This last fact is easily verified under our hypothesis that μ, ν are dominant and $\nu \geq \mu$; after all $t^\nu \geq t^\mu$, the latter equation is arrived at by right multiplication by w_0 , and t^ν is a minimal double coset representative for W in the extended affine Weyl group. □

3 Yangians

3.1 The Drinfeld Yangian

As mentioned in the introduction, we will study subquotients of Yangians in order to quantize our slices. We will actually need a slight variant on the usual Yangian, which will be produced via a theory developed by Gavarini [Ga1, Ga2]. We begin with the usual Yangian which we call the “Drinfeld Yangian” to avoid confusion with the Yangian we wish to consider.

We define the **Drinfeld Yangian** $U_h\mathfrak{g}[t]$ as the associative $\mathbb{C}[[h]]$ -algebra with generators $e_i^{(s)}, h_i^{(s)}, f_i^{(s)}$ for $i \in I$ and $r, s \in \mathbb{N}$ and relations

$$\begin{aligned} [h_i^{(s)}, h_j^{(s)}] &= 0, \\ [e_i^{(r)}, f_i^{(s)}] &= \delta_{ij} h_i^{(r+s)}, \\ [h_i^{(0)}, e_j^{(s)}] &= (\alpha_i, \alpha_j) e_j^{(s)}, \\ [h_i^{(r+1)}, e_j^{(s)}] - [h_i^{(r)}, e_j^{(s+1)}] &= \frac{h(\alpha_i, \alpha_j)}{2} (h_i^{(r)} e_j^{(s)} + e_j^{(s)} h_i^{(r)}), \\ [h_i^{(0)}, f_j^{(s)}] &= -(\alpha_i, \alpha_j) f_j^{(s)}, \\ [h_i^{(r+1)}, f_j^{(s)}] - [h_i^{(r)}, f_j^{(s+1)}] &= -\frac{h(\alpha_i, \alpha_j)}{2} (h_i^{(r)} f_j^{(s)} + f_j^{(s)} h_i^{(r)}), \\ [e_i^{(r+1)}, e_j^{(s)}] - [e_i^{(r)}, e_j^{(s+1)}] &= \frac{h(\alpha_i, \alpha_j)}{2} (e_i^{(r)} e_j^{(s)} + e_j^{(s)} e_i^{(r)}), \\ [f_i^{(r+1)}, f_j^{(s)}] - [f_i^{(r)}, f_j^{(s+1)}] &= -\frac{h(\alpha_i, \alpha_j)}{2} (f_i^{(r)} f_j^{(s)} + f_j^{(s)} f_i^{(r)}), \\ i \neq j, N = 1 - a_{ij} \Rightarrow \text{sym}[e_i^{(r_1)}, [e_i^{(r_2)}, \dots [e_i^{(r_N)}, e_j^{(s)}] \dots]] &= 0 \\ i \neq j, N = 1 - a_{ij} \Rightarrow \text{sym}[f_i^{(r_1)}, [f_i^{(r_2)}, \dots [f_i^{(r_N)}, f_j^{(s)}] \dots]] &= 0 \end{aligned}$$

where sym denotes symmetrization with respect to r_1, \dots, r_N .

The following result of Drinfeld will be our starting point.

Theorem 3.1. *$U_h\mathfrak{g}[t]$ is a quantization of $\mathfrak{g}[t]$. More precisely, there is an isomorphism of co-Poisson Hopf algebras $U_h\mathfrak{g}[t]/hU_h\mathfrak{g}[t] \cong U\mathfrak{g}[t]$, where $U\mathfrak{g}[t]$ carries the co-Poisson structure coming from the Manin triple $(\mathfrak{g}[t], t^{-1}\mathfrak{g}[[t^{-1}]], \mathfrak{g}((t^{-1})))$.*

3.2 PBW basis for the Drinfeld Yangian

Fix any order on the nodes of the Dynkin diagram; for each positive root α , we let $\check{\alpha}$ denote the smallest simple root such that $\hat{\alpha} = \alpha - \check{\alpha}$ is again a positive root.

We define $e_\alpha \in \mathfrak{g}$ for $\alpha \in \Delta_+$ recursively, by

$$e_{\alpha_i} = e_i \text{ and } e_\alpha = [e_{\hat{\alpha}}, e_{\check{\alpha}}]$$

We extend this definition to $U_h\mathfrak{g}[t]$ by defining

$$e_{\alpha_i}^{(r)} = e_i^{(r)} \text{ and } e_{\alpha}^{(r)} = [e_{\alpha}^{(r)}, e_{\alpha}^{(0)}].$$

Similarly, we define f_{α} and $f_{\alpha}^{(r)}$. We have the following PBW theorem for the Drinfeld Yangian:

Proposition 3.2.

1. Under the isomorphism $U_h\mathfrak{g}[t]/hU_h\mathfrak{g}[t] \cong U\mathfrak{g}[t]$, $e_{\alpha}^{(r)}$ corresponds to $e_{\alpha}t^r$.
2. Ordered monomials in the $e_{\alpha}^{(r)}, h_i^{(r)}, f_{\beta}^{(r)}$ form a PBW basis for $U_h\mathfrak{g}[t]$.

3.3 Drinfeld-Gavarini duality

Our goal is to give a quantization of the Poisson-Hopf algebra $\mathcal{O}(G_1[[t^{-1}]])$ using the Drinfeld Yangian $U_h\mathfrak{g}[t]$. For this we will use the quantum groups duality of Drinfeld-Gavarini.

We briefly describe one half of Drinfeld-Gavarini duality [Dr1, Ga1, Ga2]. Let (H, Δ, ϵ) be a Hopf algebra over $\mathbb{C}[[h]]$. Consider maps $\Delta^n : H \rightarrow H^{\otimes n}$ for $n \geq 0$ defined by $\Delta^0 = \epsilon$, $\Delta^1 = \text{id}_H$, and $\Delta^n = (\Delta \otimes \text{id}^{\otimes(n-2)}) \circ \Delta^{n-1}$ for $n \geq 2$. Let $\delta^n = (\text{id}_H - \epsilon)^{\otimes n} \circ \Delta^n$, and define the Hopf subalgebra

$$H' = \{a \in H \mid \delta^n(a) \in h^n H^{\otimes n}\}.$$

In general, H'/hH' is a commutative Hopf algebra over \mathbb{C} and can be given the Poisson bracket

$$\{a + hH', b + hH'\} = h^{-1}[a, b] + hH'.$$

Suppose that G is a Poisson affine algebraic group, namely the maximal spectrum of a Poisson commutative Hopf algebra $\mathcal{O}(G)$, and let $\mathfrak{g}, \mathfrak{g}^*$ be its tangent and cotangent Lie bialgebras. Let $U_h = U_h(\mathfrak{g})$ be a quantization of $U(\mathfrak{g})$.

Theorem 3.3 ([Ga1, Theorem 2.2]). *There is an isomorphism of Poisson-Hopf algebras*

$$U_h'/hU_h' \cong \mathcal{O}(G^*)$$

where G^* is a connected algebraic group with tangent Lie bialgebra \mathfrak{g}^* .

By [Ga2], for any basis $\{\overline{x}_{\alpha}\}$ of \mathfrak{g} , there exists a lift $\{x_{\alpha}\}$ in U_h such that

- $\epsilon(x_{\alpha}) = 0$,
- U_h' is generated by $\{hx_{\alpha}\}$, and
- ordered monomials in these generators span U_h' over $\mathbb{C}[[h]]$.

In particular, if $\{\bar{x}_i\}$ generates \mathfrak{g} , then $\{hx_i + hU_h'\}$ generates U_h'/hU_h' as a Poisson algebra.

To allow for easier identification of U_h'/hU_h' and $\mathcal{O}(G^*)$, we can reformulate Theorem 3.3 as follows. Consider

$$\mathcal{L} = \text{Der}(U_h'/hU_h') := \{ \varphi : U_h'/hU_h' \rightarrow \mathbb{C} \mid \varphi(ab) = \varphi(a)\epsilon(b) + \epsilon(a)\varphi(b) \}$$

with Lie bracket

$$[\varphi, \phi](a) = (\varphi \otimes \phi)(\Delta(a) - \Delta^{op}(a))$$

and cobracket

$$\delta(\varphi)(a \otimes b) = \varphi(\{a, b\})$$

This is the Lie bialgebra of the Poisson algebraic group $\text{Spec}(U_h'/hU_h')$.

The isomorphism described in Theorem 3.3 can be rephrased as follows.

Corollary 3.4. *There is an isomorphism of Lie bialgebras $\mathfrak{g}^* \cong \mathcal{L}$ defined by*

$$\bar{y} \longmapsto \left(hx + hU_h' \longmapsto \langle \bar{y}, \bar{x} \rangle \right)$$

for x a lift of $\bar{x} \in \mathfrak{g}$, extended by the Leibniz rule. This isomorphism yields a perfect Poisson–Hopf pairing $\langle \cdot, \cdot \rangle : U(\mathfrak{g}^*) \times U_h'/hU_h' \rightarrow \mathbb{C}$.

3.4 Our Yangian

We will now apply this theory to the Drinfeld Yangian $U_h\mathfrak{g}[t]$. We let $Y := (U_h\mathfrak{g}[t])'$. We will refer to Y as the Yangian from now on. Note that it is a subalgebra of the usual Yangian.

For $X = E_\alpha, H_i, F_\alpha$, and $r \geq 1$, we define $X^{(r)} = hx^{(r-1)}$. From the general remarks above these elements generate Y and monomials in these generators give a PBW basis for Y . We define a grading on Y where $X^{(r)}$ has degree r .

Theorem 3.5. *The $X^{(r)}$ generate Y subject to the relations*

$$\begin{aligned}
[H_i^{(s)}, H_j^{(s)}] &= 0, \\
[E_i^{(r)}, F_j^{(s)}] &= h\delta_{ij}H_i^{(r+s-1)}, \\
[H_i^{(1)}, E_j^{(s)}] &= h(\alpha_i, \alpha_j)E_j^{(s)}, \\
[H_i^{(r+1)}, E_j^{(s)}] - [H_i^{(r)}, E_j^{(s+1)}] &= \frac{h(\alpha_i, \alpha_j)}{2}(H_i^{(r)}E_j^{(s)} + E_j^{(s)}H_i^{(r)}), \\
[H_i^{(1)}, F_j^{(s)}] &= -h(\alpha_i, \alpha_j)F_j^{(s)}, \\
[H_i^{(r+1)}, F_j^{(s)}] - [H_i^{(r)}, F_j^{(s+1)}] &= -\frac{h(\alpha_i, \alpha_j)}{2}(H_i^{(r)}F_j^{(s)} + F_j^{(s)}H_i^{(r)}), \\
[E_i^{(r+1)}, E_j^{(s)}] - [E_i^{(r)}, E_j^{(s+1)}] &= \frac{h(\alpha_i, \alpha_j)}{2}(E_i^{(r)}E_j^{(s)} + E_j^{(s)}E_i^{(r)}), \\
[F_i^{(r+1)}, F_j^{(s)}] - [F_i^{(r)}, F_j^{(s+1)}] &= -\frac{h(\alpha_i, \alpha_j)}{2}(F_i^{(r)}F_j^{(s)} + F_j^{(s)}F_i^{(r)}), \\
i \neq j, N = 1 - a_{ij} &\Rightarrow \text{sym}[E_i^{(r_1)}, [E_i^{(r_2)}, \dots [E_i^{(r_N)}, E_j^{(s)}] \dots]] = 0 \\
i \neq j, N = 1 - a_{ij} &\Rightarrow \text{sym}[F_i^{(r_1)}, [F_i^{(r_2)}, \dots [F_i^{(r_N)}, F_j^{(s)}] \dots]] = 0 \\
E_{\alpha_i} &= E_i \\
[E_{\hat{\alpha}}^{(r)}, E_{\hat{\alpha}}^{(1)}] &= hE_{\alpha}^{(r)} \\
F_{\alpha_i} &= F_i \\
[F_{\hat{\alpha}}^{(r)}, F_{\hat{\alpha}}^{(1)}] &= hF_{\alpha}^{(r)}
\end{aligned}$$

We can repackage these generators relations using generating series. Let

$$E_i(u) = \sum_{s=1}^{\infty} E_i^{(s)} u^{-s}, \quad H_i(u) = 1 + \sum_{s=1}^{\infty} H_i^{(s)} u^{-s}, \quad F_i(u) = \sum_{s=1}^{\infty} F_i^{(s)} u^{-s}$$

Then the above relations can be written in series form. For example the series version of the commutator relation between E_i and F_i is

$$[E_i(u), F_j(v)] = \delta_{ij} \frac{h}{u-v} (H_i(u) - H_i(v)), \quad (4)$$

Remark 3.6. Note that the Drinfeld Yangian $U_h \mathfrak{g}[t]$ and our Yangian Y have natural $\mathbb{C}[h]$ -forms; moreover their $h = 1$ specializations $U_1 \mathfrak{g}[t]$ and Y_1 coincide as Hopf algebras. The gradings on $U_h \mathfrak{g}[t]$ and on Y give rise to two different filtrations on Y_1 . In the work of Brundan-Kleshchev [BrKl1], these filtrations appear as the “loop filtration” and the “Kazhdan filtration”, respectively.

3.5 Identification of Yangian with functions of $G_1[[t^{-1}]]$

From the results described above, we can deduce that there is a perfect Hopf pairing between $U(t^{-1}\mathfrak{g}[[t^{-1}]])$ and Y/hY , as per Corollary 3.4. Let us denote by Q the root lattice for \mathfrak{g} , let Q_+ denote the positive root cone, and let $Q_{>} = Q_+ \setminus \{0\}$, $Q_{<} = -Q_{>}$.

Lemma 3.7. *The Drinfeld Yangian $U_h\mathfrak{g}[t]$, Y , and Y/hY , are all Q -graded Hopf algebras (all tensor products being graded by total degree). The pairing between $U(t^{-1}\mathfrak{g}[[t^{-1}]])$ and Y/hY respects this grading.*

Proof. The Hopf grading on these spaces is induced by the action of the elements $h_i^{(0)}$ (resp. $H_i^{(1)}$). In each case, coproducts preserve total degree since the coproduct is a homomorphism and the above elements are Lie algebra-like.

It is clear from the formulas of Corollary 3.4 that the pairing between $U(t^{-1}\mathfrak{g}[[t^{-1}]])$ and Y/hY respects the grading for pairings $\langle y, x \rangle$, when $y \in t^{-1}\mathfrak{g}[[t^{-1}]]$, $x \in Y_0$. The result follows for monomials $y_1 \cdots y_k \in U(t^{-1}\mathfrak{g}[[t^{-1}]])$ by induction on k . \square

For $\alpha \in Q$, let $Y(\alpha)$ be the corresponding component of Y/hY as per Lemma 3.7.

Proposition 3.8. *In Y/hY we have:*

$$\begin{aligned} \Delta(H_i^{(r)}) &= H_i^{(r)} \otimes 1 + 1 \otimes H_i^{(r)} + \sum_{s=1}^{r-1} H_i^{(s)} \otimes H_i^{(r-s)} + \bigoplus_{\substack{\alpha+\beta=0 \\ \alpha \in Q_{<}, \beta \in Q_{>}}} Y(\alpha) \otimes Y(\beta) \\ \Delta(E_i^{(r)}) &= E_i^{(r)} \otimes 1 + 1 \otimes E_i^{(r)} + \sum_{s=1}^{r-1} H_i^{(s)} \otimes E_i^{(r-s)} + \bigoplus_{\substack{\alpha+\beta=\alpha_i \\ \alpha \in Q_{<}, \beta \in Q_{>}}} Y(\alpha) \otimes Y(\beta) \\ \Delta(F_i^{(r)}) &= F_i^{(r)} \otimes 1 + 1 \otimes F_i^{(r)} + \sum_{s=1}^{r-1} F_i^{(s)} \otimes H_i^{(r-s)} + \bigoplus_{\substack{\alpha+\beta=-\alpha_i \\ \alpha \in Q_{<}, \beta \in Q_{>}}} Y(\alpha) \otimes Y(\beta) \end{aligned}$$

Proof. To begin we recall that $\Delta(X^{(1)}) = X^{(1)} \otimes 1 + 1 \otimes X^{(1)}$ for all $x \in \mathfrak{g}$. Also, using the presentation of $U_h\mathfrak{g}[t]$ with generators $x, J(x)$ for $x \in \mathfrak{g}$ (for which the coproduct is known), a direct calculation yields

$$\Delta(H_i^{(2)}) = H_i^{(2)} \otimes 1 + 1 \otimes H_i^{(2)} + H_i^{(1)} \otimes H_i^{(1)} - \sum_{\beta \in \Phi_+} C_\beta(\beta, \alpha_i) F_\beta^{(1)} \otimes E_\beta^{(1)}$$

where $(e_\beta, f_\beta) = C_\beta^{-1}$. We prove the coproduct for $E_i^{(r)}$ by induction on r , using the identity

$$E_i^{(r+1)} = \frac{1}{(\alpha_i, \alpha_i)} \left\{ H_i^{(2)}, E_i^{(r)} \right\} - H_i^{(1)} E_i^{(r)}$$

The coproduct of the right side is expanded using the Poisson-Hopf algebra relations, the formula for $\Delta(H_i^{(2)})$, and the inductive hypothesis. The above identity is then applied again to reduce the terms in the result, and yields the form as claimed.

An analogous induction proves the case of $\Delta(F_i^{(r)})$. Finally, we take the coproduct of the identity

$$H_i^{(r)} = \{E_i^{(r)}, F_i^{(1)}\}$$

to finish the proof. \square

Recall that the pairing between $U(t^{-1}\mathfrak{g}[[t^{-1}]])$ and Y/hY is determined, as per Corollary 3.4, by the pairing between $t^{-1}\mathfrak{g}[[t^{-1}]]$ and $\mathfrak{g}[t]$ given in Section 2.6. Choose an ‘‘FHE’’ total ordering on the generators $f_\alpha t^r, h_i t^r, e_\alpha t^r$ for $U(t^{-1}\mathfrak{g}[[t^{-1}]])$. Then it is easy to see that the previous lemma and proposition completely control the pairing between $U(t^{-1}\mathfrak{g}[[t^{-1}]])$ and Y/hY for the corresponding PBW basis. For example, $-F_i^{(r)}$ acts as the dual of the basis element $e_i t^{-r}$, etc.

Theorem 3.9. *There is an isomorphism of \mathbb{N} -graded Poisson Hopf algebras $\phi: Y/hY \cong \mathcal{O}(G_1[[t^{-1}]])$ such that*

$$\begin{aligned}\phi(H_i(u)) &= \prod_j \Delta_{\omega_j, \omega_j}(u)^{-a_{ji}} \\ \phi(F_i(u)) &= d_i^{-1/2} \frac{\Delta_{\omega_i, s_i \omega_i}(u)}{\Delta_{\omega_i, \omega_i}(u)} \\ \phi(E_i(u)) &= d_i^{-1/2} \frac{\Delta_{s_i \omega_i, \omega_i}(u)}{\Delta_{\omega_i, \omega_i}(u)}\end{aligned}$$

where $\mathcal{O}(G_1[[t^{-1}]])$ is graded using the loop rotation \mathbb{C}^\times action.

Proof. We check explicitly that the right-hand sides act as described by the previous proposition. Let $X = (x_1 t^{r_1}) \cdots (x_k t^{r_k}) \in U(t^{-1}\mathfrak{g}[[t^{-1}]])$ be a basis monomial with the FHE order as chosen above. Then we have

$$\frac{\Delta_{\omega_i, s_i \omega_i}(u)}{\Delta_{\omega_i, \omega_i}(u)}(X) = -d_i^{-1/2} \frac{\partial^k}{\partial z_1 \cdots \partial z_k} \frac{\langle v_{-\omega_i}, (1 + z_1 u^{r_1} x_1) \cdots (1 + z_k u^{r_k} x_k) f_i v_{\omega_i} \rangle}{\langle v_{-\omega_i}, (1 + z_1 u^{r_1} x_1) \cdots (1 + z_k u^{r_k} x_k) v_{\omega_i} \rangle} \Big|_{z_1 = \dots = z_k = 0}$$

noting that $\overline{s_i} v_{\omega_i} = f'_i v_{\omega_i} = -d_i^{-1/2} f_i v_{\omega_i}$ in the generalized minor (see Section 2.1). Since we have an FHE order, to get something nonzero in the right-hand numerator x_k must be a multiple of e_i , since $e_i f_i v_{\omega_i} = h_i v_{\omega_i} = d_i v_{\omega_i}$. In this case $z_k u^{r_k} e_i$ does not contribute to the denominator, and the remaining factors cancel leaving

$$\frac{\Delta_{\omega_i, s_i \omega_i}(u)}{\Delta_{\omega_i, \omega_i}(u)}(X) = -d_i^{1/2} \frac{\partial^k}{\partial z_1 \cdots \partial z_k} z_k u^{r_k} \Big|_{z_1 = \dots = z_k = 0}$$

so X must have been $e_i t^r$ to start with. But this is precisely how $d_i^{1/2} F_i(u)$ acts on X . Similar computations hold in the two remaining cases.

To prove the equality for $H_i(u)$ one can also work in $\mathcal{O}(G_1[[t^{-1}]])$, and build off the known results $E_i(u)$ and $F_i(u)$, since we must have

$$\frac{\phi(H_i(u)) - \phi(H_i(v))}{u - v} = \left\{ \phi(E_i(u)), \phi(F_i(v)) \right\}$$

We can then use formula (1) and identities for generalized minors.

The nondegeneracy of both Hopf pairings implies that ϕ is an injection. It follows that ϕ is an isomorphism from a dimension count; both Y/hY and $\mathcal{O}(G_1[[t^{-1}]])$ have Hilbert series for the loop grading given by

$$\prod_{i=1}^{\infty} \frac{1}{(1 - q^i)^{\dim \mathfrak{g}}}.$$

Indeed for Y/hY this follows from the PBW theorem coming from Y , since Y is a free $\mathbb{C}[[h]]$ -algebra. On the other hand, the Hilbert series on $\mathcal{O}(G_1[[t^{-1}]])$ is the same as the Hilbert series for $\text{Sym}(t^{-1} \mathfrak{g}[[t^{-1}]])$ since as $G_1[[t^{-1}]]$ is pro-unipotent, we have an isomorphism of vector spaces.

□

3.6 Shifted Yangians

The Yangian has a very interesting class of subalgebras: the shifted Yangians. Let μ be a dominant weight.

We will now redefine elements

$$F_{\alpha}^{(s)} = \frac{1}{h} [F_{\check{\alpha}}^{(s - \langle \mu^*, \check{\alpha} \rangle)}, F_{\check{\alpha}}^{(\langle \mu^*, \check{\alpha} \rangle + 1)}]. \quad (5)$$

for α a positive non-simple root and for $s > \langle \mu^*, \alpha \rangle$. Note that these $F_{\alpha}^{(s)}$ depend on μ .

Definition 3.10. The **shifted Yangian** Y_{μ} is the subalgebra of Y generated by $E_{\alpha}^{(s)}$ for all α, s , $H_i^{(s)}$ for all i, s , and $F_{\alpha}^{(s)}$ for $s > \langle \mu^*, \alpha \rangle$.

Proposition 3.11. 1. *Monomials in the $E_{\alpha}^{(s)}, H_i^{(s)}, F_{\alpha}^{(s)}$ give a basis for Y_{μ} .*

2. *The natural map $Y_{\mu}/hY_{\mu} \rightarrow Y/hY$ is injective.*

Proof. We first construct a PBW basis for Y slightly different from the one described in Section 3.4. The generators $E_{\alpha}^{(s)}$ are defined as usual (cf. Section 3.4). The generators $F_{\alpha}^{(s)}$ are given the usual definition when $s \leq \langle \mu^*, \alpha \rangle$, but for

$s > \langle \mu^*, \alpha \rangle$ we take definition (5). By the general remarks following Theorem 3.3, ordered monomials in generators $F_\alpha^{(s)}, H_i^{(s)}, E_\alpha^{(s)}$ are a PBW basis of Y .

Any element $x \in Y_\mu$ can be expressed as a linear combination of these PBW monomials. We now show that any monomials appearing in such an expression do not contain factors of the form $F_\alpha^{(s)}$ for $s \leq \langle \mu^*, \alpha \rangle$.

By definition, x is a linear combination of (unordered) monomials in $F_\alpha^{(s)}, H_i^{(t)}, E_\alpha^{(u)}$, where $s > \langle \mu^*, \alpha \rangle$. To put x in PBW form one has to commute these generators past each other. By definition, when $s > \langle \mu^*, \alpha \rangle$, $F_\alpha^{(s)}$ is a linear combination of monomials built from $F_i^{(t)}$, where $t > \langle \mu^*, \alpha_i \rangle$. Therefore it suffices to show that when commuting such $F_i^{(t)}$ past the other generators of Y_μ one never obtains factors of the form $F_j^{(u)}$ for $u \leq \langle \mu^*, \alpha_j \rangle$. This is a direct consequence of the relations appearing in Theorem 3.5.

This proves the first statement of the theorem. The second part is a direct consequence of the first. \square

In the limit as $\mu \rightarrow \infty$, then we obtain Y_∞ which is the subalgebra generated by all $E_\alpha^{(s)}, A_i^{(s)}$. This is called the **Borel Yangian** in [FR].

We will now show that this shifted Yangian is a quantization of Gr_μ . Recall that $\mathcal{O}(\text{Gr}_\mu)$ is embedded as a Poisson subalgebra of $\mathcal{O}(G_1[[t^{-1}]])$.

Theorem 3.12. *The isomorphism ϕ restricts to an isomorphism of Poisson algebras from Y_μ/hY_μ to $\mathcal{O}(\text{Gr}_\mu)$.*

Proof. First note that Y_μ/hY_μ is generated as a Poisson algebra by all $E_i^{(s)}, A_i^{(s)}$, and those $F_i^{(s)}$ for $i > \langle \mu^*, \alpha_i \rangle$. We note that Lemma 2.19 shows that the image of these generators under ϕ land in the subalgebra $\mathcal{O}(\text{Gr}_\mu)$.

Since $\mathcal{O}(\text{Gr}_\mu)$ is a Poisson subalgebra of $\mathcal{O}(G_1[[t^{-1}]])$, we see that ϕ restricts to a map $Y_\mu/hY_\mu \rightarrow \mathcal{O}(G_1[[t^{-1}]])$. This map is injective, since it is the restriction of an injective map. Thus, we only need to show that it is surjective, which we do by a dimension count.

Note that by Lemma 2.3 the isotropy Lie algebra of $t^{w_0\mu}$ in $G_1[[t^{-1}]]$ is the finite dimensional nilpotent Lie algebra

$$\bigoplus_{\alpha \in \Delta_+} \bigoplus_{i=1}^{-\langle w_0\mu, \alpha \rangle} t^{-i} \mathfrak{g}_\alpha.$$

As a \mathbb{C}^* -module, the functions on the group are identical to those on the Lie algebra by the unipotence of the stabilizer. Thus, if we let $d(k)$ be the number of roots such that $\langle w_0\mu, \alpha \rangle < k$, the Hilbert series of the functions on the stabilizer

is

$$\prod_{i=1}^{\infty} \frac{1}{(1 - q^i)^{d(i)}}.$$

The Hilbert series of $\mathcal{O}(G_1[[t^{-1}]])^{G_1[[t^{-1}]]_\mu}$ is the quotient of that of $\mathcal{O}(G_1[[t^{-1}]])$ by that of functions on the stabilizer. That is, it is

$$\prod_{i=1}^{\infty} \frac{1}{(1 - q^i)^{\dim \mathfrak{g} - d(i)}}.$$

On the other hand, the PBW basis for the shifted Yangian gives us the same Hilbert series for Y_μ . \square

Thus the shifted Yangian Y_μ gives a quantization of \mathbf{Gr}_μ .

Remark 3.13. We should note that it is this theorem that forces us to use the thick Grassmannian; it will fail if we take the analogue of \mathbf{Gr}_μ in the thin affine Grassmannian, since this has “too many” functions, and will correspond to a completion of Y_μ .

3.7 Deformation of the Yangian

We consider a deformation of the Yangian, which we think of as related to the Beilinson-Drinfeld Grassmannian deforming the affine Grassmannian. We consider for each node i in the Dynkin diagram an infinite sequence of parameters $r_i^{(1)}, r_i^{(2)}, \dots \in \mathbb{C}[[h]]$ and their generating series $r_i(u) = 1 + r_i^{(1)}u^{-1} + \dots$.

Now consider the algebra $Y(\mathbf{r})$ generated by the coefficients of $E_i(u), F_i(u), A_i(u)$. The relations are as in the previous section, with the relation (4) replaced by

$$(u - v)[E_i(u), F_i(v)] = h(r_i(u)H_i(u) - r_i(v)H_i(v)), \quad (6)$$

and let $Y_\mu(\mathbf{r})$ be the shifted analogue of this algebra. $Y(\mathbf{r})$ is actually isomorphic to the trivial deformation of the Yangian via the map $H_i(u) \mapsto H_i(u)/r_i(u)$.

4 Quantization of slices

In order to quantize the slices \mathbf{Gr}_μ^λ , we will need to define a quotient of Y_μ (and its deformations $Y_\mu(\mathbf{r})$). To do this we will use the work of Gerasimov-Kharchev-Lebedev-Oblezin [GKLO].

4.1 Change of Cartan generators

It will be convenient for us to change the Cartan generators of Y . Following [GKLO], we define $A_i^{(s)}$ by the equation

$$H_i(u) = \frac{\prod_{j \neq i} \prod_{p=1}^{-a_{ji}} A_j(u - \frac{h}{2}(\alpha_i + p\alpha_j, \alpha_j))}{A_i(u)A_i(u - \frac{h}{2}(\alpha_i, \alpha_i))} \quad (7)$$

where $A_i(u) = 1 + \sum_{s=1}^{\infty} A_i^{(s)} u^{-s}$.

Example 4.1. In the $G = SL_2$ case, this gives $H(u) = \frac{1}{A(u)A(u-h)}$ and so for example we have

$$H^{(1)} = -2A^{(1)}, \quad H^{(2)} = 3A^{(1)^2} - hA^{(1)} - 2A^{(2)}$$

Proposition 4.2 ([GKLO, Lemma 2.1]). *Equation 7 uniquely determines all the $A_i^{(s)}$.* \square

One can think of the new generators $A_i^{(s)}$ as being related to the fundamental coweights of G , whereas the $H_i^{(s)}$ match with the simple coroots. In particular, we have the following result which follows by setting $h = 0$ in (7).

Proposition 4.3. *Let $\phi : Y/hY \rightarrow \mathcal{O}(G_1[[t^{-1}]])$ be the isomorphism from Theorem 3.9. Then $\phi(A_i^{(s)}) = \Delta_{\omega_i, \omega_i}^{(s)}$.* \square

4.2 The GKLO representation

In this section, we describe certain representations via difference operators of shifted Yangians, based on work of Gerasimov-Kharchev-Lebedev-Oblezin [GKLO]. Fix an orientation of the Dynkin diagram; we will write $i \leftarrow j$ to denote arrows in this quiver. This will replace the ordering on the simple roots in [GKLO].

Fix a dominant weight λ such that $\mu \leq \lambda$ and let $m_i = \langle \lambda - \mu, \omega_{i^*} \rangle$ and let $\lambda_i = \langle \lambda, \alpha_{i^*} \rangle$.

Define a $\mathbb{C}[[h]]$ -algebra D_μ^λ , with generators $z_{i,k}, \beta_{i,k}, \beta_{i,k}^{-1}$, for $i \in I$ and $1 \leq k \leq m_i$, and $(z_{i,k} - z_{i,l})^{-1}$, and relations that all generators commute except that $\beta_{i,k} z_{i,k} = (z_{i,k} + d_i h) \beta_{i,k}$.

This algebra D_μ^λ is an algebra of h -difference operators.

Proposition 4.4. *The algebra D_μ^λ is a free $\mathbb{C}[[h]]$ -algebra and we have an isomorphism of Poisson algebras*

$$D_\mu^\lambda / h D_\mu^\lambda \cong \mathbb{C}[z_{i,k}, (z_{i,k} - z_{i,l})^{-1}, \beta_{i,k}, \beta_{i,k}^{-1}]$$

where the right hand side is given the Poisson structure defined by $\{\beta_{i,k}, z_{i,k}\} = d_i \beta_{i,k}$ and all other generators Poisson commute.

Proof. Obviously, we have a map

$$\mathbb{C}[z_{i,k}, (z_{i,k} - z_{i,l})^{-1}, \beta_{i,k}, \beta_{i,k}^{-1}] \rightarrow D_\mu^\lambda / hD_\mu^\lambda$$

by observing that $D_\mu^\lambda / hD_\mu^\lambda$ is commutative. From the Bergman diamond lemma, we see that the algebra D_μ^λ has a PBW basis consisting of

$$h^p \cdot \prod \beta_{i,k}^{\pm a_{i,k}} \cdot \prod z_{j,k}^{b_{j,k}} \cdot \prod_{k < \ell} (z_{i,k} - z_{i,\ell})^{e_{i,k,\ell}}$$

subject to restriction that if $b_{j,k} \neq 0$, then k must be maximal in its equivalence class for the relation given by the transitive closure of the binary relation $k \sim \ell$ if $e_{j,k,\ell} \neq 0$. Freeness over $\mathbb{C}[[h]]$ follows immediately and since the same monomials give a basis of $\mathbb{C}[[h]][z_{i,k}, (z_{i,k} - z_{i,l})^{-1}, \beta_{i,k}, \beta_{i,k}^{-1}]$, this confirms that we have the desired isomorphism. The Poisson bracket calculation follows immediately from the relations. \square

Fix some complex numbers $c_i^{(r)}$ for $i \in I$, $1 \leq r \leq \lambda_i$. For any variable x , consider the monic degree λ_i polynomial whose coefficients are the numbers $c_i^{(r)}$, $C_i(x) = x^{\lambda_i} + c_i^{(1)} x^{\lambda_i-1} + \dots + c_i^{(\lambda_i)}$. Note that $x^{-\lambda_i} C_i(x) = 1 + c_i^{(1)} x^{-1} + \dots + c_i^{(\lambda_i)} x^{-\lambda_i}$. We also introduce polynomials $Z_i(x) = \prod_{k=1}^{m_i} (x - z_{i,k})$ and $Z_{i,k}(x) = \prod_{\ell \neq k} (x - z_{i,\ell})$. Let $\mu_i = \langle \mu, \alpha_{i^*} \rangle$ and set $F_{\mu,i}(u) = \sum_{s=1}^{\infty} F_i^{(s+\mu_i)} u^{-s}$. Finally, for any \mathbf{c} as above, define \mathbf{r} by

$$r_i(u) = u^{-\lambda_i} C_i(u) \frac{\prod_{j \neq i} \prod_{p=1}^{-a_{ji}} (1 - u^{-1}(hd_i \frac{a_{ij}}{2} + hd_j p))^{m_j}}{(1 - hd_i u^{-1})^{m_i}} \quad (8)$$

We are now ready to define the **GKLO representation**:

Theorem 4.5. *There is a map of $\mathbb{C}[[h]]$ -algebras, $\Psi_\mu^\lambda : Y_\mu(\mathbf{r}) \rightarrow D_\mu^\lambda$ defined by:*

$$\begin{aligned} A_i(u) &\mapsto u^{-m_i} Z_i(u) \\ E_i(u) &\mapsto d_i^{-1/2} \sum_{k=1}^{m_i} \frac{\prod_{j \rightarrow i} \prod_{p=1}^{-a_{ji}} Z_j(z_{i,k} - hd_i \frac{a_{ij}}{2} - hd_j p)}{(u - z_{i,k}) Z_{i,k}(z_{i,k})} \beta_{i,k}^{-1} \end{aligned}$$

And, $F_{\mu,i}(u)$ maps to

$$-d_i^{-1/2} \sum_{k=1}^{m_i} C_i(z_{i,k} + hd_i) \frac{\prod_{j \leftarrow i} \prod_{p=1}^{-a_{ji}} Z_j(z_{i,k} - hd_i(\frac{a_{ij}}{2} - 1) - hd_j p)}{(u - z_{i,k} - hd_i) Z_{i,k}(z_{i,k})} \beta_{i,k}$$

Proof. When $\mu = 0$ this is a reformulation of [GKLO, Theorem 3.1.(i)]. Suppose then that $\mu \neq 0$. Then the proof of [GKLO, Theorem 3.1] applies to all the relations in Y_μ except for the commutator relation between $E_i(u)$ and $F_{\mu,i}(v)$.

In the shifted Yangian this relation takes the form

$$(u - v)[E_i(u), F_{\mu,i}(v)] = h(J_{\mu,i}(v) - J_{\mu,i}(u)) \quad (9)$$

where $J_i(v) = r_i(v)H_i(v) = \sum_{p=0}^{\infty} J_i^{(p)}v^{-p}$ and

$$J_{\mu,i}(v) = \sum_{p=1}^{\infty} J_i^{(p+\mu_i)}v^{-p}.$$

To express the left hand side of (9) we set:

$$\begin{aligned} L_i(v) &= \frac{C_i(z_{i,k} + hd_i) \prod_{j \neq i} \prod_{p=1}^{-a_{ji}} Z_j(z_{i,k} - hd_i(\frac{a_{ij}}{2} - 1) - hd_j p)}{Z_{i,k}(z_{i,k} + hd_i) Z_{i,k}(z_{i,k})(v - z_{i,k} - hd_i)} \\ R_i(v) &= \frac{C_i(z_{i,k}) \prod_{j \neq i} \prod_{p=1}^{-a_{ji}} Z_j(z_{i,k} - hd_i(\frac{a_{ij}}{2} - hd_j p))}{Z_{i,k}(z_{i,k} - hd_i) Z_{i,k}(z_{i,k})(v - z_{i,k})} \end{aligned}$$

Then the left hand side of (9) is equal to

$$d_i^{-1} \sum_{k=1}^{m_i} (L_i(v) - R_i(v)) - (L_i(u) - R_i(u))$$

Note that we expressed this sum as a “ v -part” minus a “ u -part”.

Now we consider the right hand side of (9). Note that

$$\lambda_i = \mu_i + 2m_i + \sum_{j \leftrightarrow i} a_{ji} m_j$$

Therefore,

$$r_i(u) = u^{-\mu_i} \frac{C_i(u) \prod_{j \neq i} \prod_{p=1}^{-a_{ji}} (u - hd_i(\frac{a_{ij}}{2} - hd_j p))^{m_j}}{u^{m_i} (u - hd_i)^{m_i}}$$

Now

$$H_i(u) \mapsto \frac{u^{m_i} (u - hd_i)^{m_i}}{\prod_{j \neq i} \prod_{p=1}^{-a_{ji}} (u - hd_i(\frac{a_{ij}}{2} - hd_j p))^{m_j}} \frac{\prod_{j \neq i} \prod_{p=1}^{-a_{ji}} Z_j(u - hd_i(\frac{a_{ij}}{2} - hd_j p))}{Z_i(u) Z_i(u - hd_i)}$$

and hence

$$r_i(u) H_i(u) \mapsto u^{-\mu_i} C_i(u) \frac{\prod_{j \neq i} \prod_{p=1}^{-a_{ji}} Z_j(u - hd_i(\frac{a_{ij}}{2} - hd_j p))}{Z_i(u) Z_i(u - hd_i)}$$

Therefore

$$C_i(u) \frac{\prod_{j \neq i} \prod_{p=1}^{-a_{ji}} Z_j(u - hd_i \frac{a_{ij}}{2} - hd_j p)}{Z_i(u) Z_i(u - hd_i)} = \sum_{p=0}^{\infty} J_i^{(p)} u^{\mu_i - p}$$

On the other hand

$$J_{\mu,i}(u) = \sum_{p=\mu_i+1}^{\infty} J_i^{(p)} u^{\mu_i - p}$$

showing that $J_{\mu,i}(u)$ is a truncation of $C_i(u) \frac{\prod_{j \neq i} \prod_{p=1}^{-a_{ji}} Z_j(u - hd_i \frac{a_{ij}}{2} - hd_j p)}{Z_i(u) Z_i(u - hd_i)}$. More precisely, for $r = 1, 2, \dots$

$$hJ_{\mu,i}(u) \Big|_{u^{-r}} = hC_i(u) \frac{\prod_{j \neq i} \prod_{p=1}^{-a_{ji}} Z_j(u - hd_i \frac{a_{ij}}{2} - hd_j p)}{Z_i(u) Z_i(u - hd_i)} \Big|_{u^{-r}}$$

Using partial fractions we have that $\frac{h}{Z_i(u) Z_i(u - hd_i)}$ equals

$$\sum_{k=1}^{m_i} \frac{1}{Z_{ik}(z_{ik}) Z_{ik}(z_{ik} + hd_i) (u - z_{ik} - hd_i)} - \frac{1}{Z_{ik}(z_{ik}) Z_{ik}(z_{ik} - hd_i) (u - z_{ik})}$$

Therefore for $r = 1, 2, \dots$ the u^{-r} -coefficient of $hJ_{\mu,i}(u)$ is equal to the u^{-r} -coefficient of

$$\begin{aligned} & \sum_{k=1}^{m_i} \frac{C_i(u) \prod_{j \neq i} \prod_{p=1}^{-a_{ji}} Z_j(u - hd_i \frac{a_{ij}}{2} - hd_j p)}{Z_{ik}(z_{ik}) Z_{ik}(z_{ik} + hd_i) (u - z_{ik} - hd_i)} - \\ & \frac{C_i(u) \prod_{j \neq i} \prod_{p=1}^{-a_{ji}} Z_j(u - hd_i \frac{a_{ij}}{2} - hd_j p)}{Z_{ik}(z_{ik}) Z_{ik}(z_{ik} - hd_i) (u - z_{ik})} \end{aligned}$$

Now observe that for any polynomial $p(u)$ and for $r = 1, 2, \dots$

$$\frac{p(u)}{u - z} \Big|_{u^{-r}} = \frac{p(z)}{u - z} \Big|_{u^{-r}}$$

Therefore for $r = 1, 2, \dots$ the u^{-r} -coefficient of $hu^{\mu_i} J_{\mu,i}(u)$ is equal to the u^{-r} -coefficient of

$$\sum_{k=1}^{m_i} L_i(u) - R_i(u)$$

proving (9). □

Example 4.6. If $\mathfrak{g} = \mathfrak{sl}_2$ and $\lambda = \alpha^\vee, \mu = 0$, then the formulas above simplify considerably. In this case,

$$A(u) \mapsto 1 - zu^{-1} \quad E(u) \mapsto \frac{1}{u-z}\beta^{-1}$$

and $F(u) \mapsto -((z+h)^2 + c^{(1)}(z+h) + c^{(2)})\frac{1}{u-z-h}\beta$. In particular,

$$H^{(1)} \mapsto 2z \quad E^{(1)} \mapsto \beta^{-1} \quad F^{(1)} \mapsto -((z+h)^2 + c^{(1)}(z+h) + c^{(2)})\beta$$

Restrict this representation to the copy of \mathfrak{sl}_2 generated by $E^{(1)}, H^{(1)} + c^{(1)} + h, F^{(1)}$, and consider these as difference operators acting on the polynomial ring $\mathbb{C}[z]$. (More precisely, these act on $\mathbb{C}[[h]][z]$, but one can specialize h to 1.) This is a standard Whittaker module for \mathfrak{sl}_2 with generic nilpotent character.

Remark 4.7. We can define a \mathbb{Z} -grading on D_μ^λ by setting

$$\deg h = 1, \deg z_{i,k} = 1, \deg \beta_{i,k} = m_i + \sum_{i \rightarrow j} a_{ij}m_j + \lambda_i - \mu_i$$

With this definition, the GKLO representation preserves grading.

4.3 Quantization of the slices Gr_μ^λ

For any \mathbf{c} as above, let $Y_\mu^\lambda(\mathbf{c})$ be the image of $Y_\mu(\mathbf{r})$ in D_μ^λ under the GKLO representation Ψ_μ^λ and let $I_\mu^\lambda(\mathbf{c})$ denote the kernel of Ψ_μ^λ (here \mathbf{r} is determined from \mathbf{c} by (8)).

Note that $Y_\mu^\lambda(\mathbf{c})$ is free as a $\mathbb{C}[[h]]$ -algebra since it is a subalgebra of D_μ^λ , a free $\mathbb{C}[[h]]$ -algebra.

We have the isomorphism $Y_\mu(\mathbf{c}) \rightarrow Y_\mu$ from section 3.7 and thus we get an isomorphism of Poisson algebras $Y_\mu(\mathbf{c})/hY_\mu(\mathbf{c}) \rightarrow \mathcal{O}(\text{Gr}_\mu)$ from Theorem 3.12. On the other hand, because $Y_\mu^\lambda(\mathbf{c})$ is free as a $\mathbb{C}[[h]]$ -algebra, we get a surjection of Poisson algebras $Y_\mu(\mathbf{c})/hY_\mu(\mathbf{c}) \rightarrow Y_\mu^\lambda(\mathbf{c})/hY_\mu^\lambda(\mathbf{c})$.

We will now establish the following theorem which shows that Y_μ^λ is a quantization of scheme supported on Gr_μ^λ .

Theorem 4.8. *There is a surjective map of Poisson algebras $Y_\mu^\lambda(\mathbf{c})/hY_\mu^\lambda(\mathbf{c}) \rightarrow \mathcal{O}(\text{Gr}_\mu^\lambda)$ which is an isomorphism modulo the nilradical of the left hand side.*

Remark 4.9. Consider the map

$$Y_\mu^\lambda(\mathbf{c})/hY_\mu^\lambda(\mathbf{c}) \rightarrow \mathbb{C}[z_{i,k}, (z_{i,k} - z_{i,l})^{-1}, \beta_{i,k}, \beta_{i,k}^{-1}]$$

obtained by reducing the GKLO representation mod h . If we knew that this map was injective, then we would know that $Y_\mu^\lambda(\mathbf{c})/hY_\mu^\lambda(\mathbf{c})$ was reduced and that the map from Theorem 4.8 was an isomorphism. We will in fact make a stronger conjecture.

If Conjecture 2.20 holds, then we can strengthen Theorem 4.8 as follows.

Theorem 4.10. *If Conjecture 2.20 holds then*

1. *There is an isomorphism of Poisson algebras $Y_\mu^\lambda(\mathbf{c})/hY_\mu^\lambda(\mathbf{c}) \rightarrow \mathcal{O}(\text{Gr}_\mu^\lambda)$.*
2. *$Y_\mu^\lambda(\mathbf{c})$ is the quotient of $Y_\mu(\mathbf{c})$ by the 2-sided ideal generated by $A_i^{(s)}$ for $s > m_i, i \in I$.*

Proof of Theorem 4.8. Via the isomorphism $Y_\mu(\mathbf{c})/hY_\mu(\mathbf{c}) \rightarrow \mathcal{O}(\text{Gr}_\mu)$, we can regard $Y_\mu^\lambda(\mathbf{c})$ as a quotient of $\mathcal{O}(\text{Gr}_\mu)$ by an ideal, which we denote I_2 .

First, note that $\Psi_\mu^\lambda(A_i^{(s)}) = 0$ for $i \in I, s > m_i$ and thus $\Delta_{\omega_i, \omega_i}^{(s)} \in I_2$ for $i \in I$ and $s > m_i$. Since I_2 is a Poisson ideal, we see that $J_\mu^\lambda \subset I_2$.

By Proposition 2.21, we see that the vanishing locus of J_μ^λ is Gr_μ^λ and thus the vanishing locus of I_2 is contained in Gr_μ^λ . Thus it suffices to show that the vanishing locus of I_2 is not strictly contained in Gr_μ^λ .

Since I_2 is a Poisson ideal, we see that $V(I_2)$ is a Poisson subvariety of Gr_μ^λ and thus is the union of $\text{Gr}_\mu^{\bar{\nu}}$, for $\nu \leq \lambda$. Suppose that we have $V(I_2) = \cup_j \text{Gr}_\mu^{\bar{\nu}_j}$ for $\nu_j < \lambda$. For each j , there exists i such that $\langle \nu_j - \mu, \omega_{i^*} \rangle < \langle \lambda - \mu, \omega_{i^*} \rangle = m_i$. Thus applying Proposition 2.4, $\prod_i \Delta_{\omega_i, \omega_i}^{(m_i)}$ vanishes on $\cup_j \text{Gr}_\mu^{\bar{\nu}_j}$. Hence for some k we have $(\prod_i \Delta_{\omega_i, \omega_i}^{(m_i)})^k \in I_2$.

On the other hand, we see that under the GKLO representation

$$\Psi_\mu^\lambda(A_i^{(m_i)}) = (-1)^{m_i} z_{i,1} \cdots z_{i,m_i}$$

and thus under the map

$$\mathcal{O}(\text{Gr}_\mu) \cong Y_\mu(\mathbf{c})/hY_\mu(\mathbf{c}) \rightarrow D_\mu^\lambda/hD_\mu^\lambda \cong \mathbb{C}[z_{i,k}, (z_{i,k} - z_{i,l})^{-1}, \beta_{i,k}, \beta_{i,k}^{-1}]$$

we see that $(\prod_i \Delta_{\omega_i, \omega_i}^{(m_i)})^k$ is mapped to a monomial in the $z_{i,k}$. In particular, this shows that $(\prod_i \Delta_{\omega_i, \omega_i}^{(m_i)})^k$ does not lie in I_2 , contradicting the previous paragraph.

Thus we conclude that $V(I_2) = \text{Gr}_\mu^\lambda$ as desired. \square

Proof of Theorem 4.10. Let I_1 be the ideal of Gr_μ^λ in $\mathcal{O}(\text{Gr}_\mu)$.

Let K be the ideal in $Y_\mu(\mathbf{c})$ generated by $A_i^{(s)}$ for $s > m_i, i \in I$. Then we have an inclusion $K \subset I_\mu^\lambda(\mathbf{c})$ and a resulting map

$$K/hK \rightarrow I_\mu^\lambda(\mathbf{c})/hI_\mu^\lambda(\mathbf{c}) = I_2$$

which may not be injective. Let I_3 denote the image of this map. From the definitions, we see that $I_3 \subset I_2$. Moreover, we have that $J_\mu^\lambda \subset I_3$, since I_3 is a Poisson ideal and it contains the generators of I_3 .

In the previous proof we have shown that $I_2 \subset I_1$. Thus we have a chain of inclusions $J_\mu^\lambda \subset I_3 \subset I_2 \subset I_1$. On the other hand, Conjecture 2.20 shows us that $I_1 = J_\mu^\lambda$.

Hence we conclude that $I_1 = I_2 = I_3 = J_\mu^\lambda$. So the first assertion holds.

For the second assertion, note that $I_3 = I_2$ implies that $K/hK \rightarrow I_\mu^\lambda(\mathbf{c})/hI_\mu^\lambda(\mathbf{c})$ is surjective. Let $L = I_\mu^\lambda(\mathbf{c})/K$. The long exact sequence for $\otimes_{\mathbb{C}[[h]]}\mathbb{C}$ gives

$$K/hK \rightarrow I_\mu^\lambda(\mathbf{c})/hI_\mu^\lambda(\mathbf{c}) \rightarrow L/hL \rightarrow 0$$

and thus $L/hL = 0$. From Nakayama's lemma, we conclude that $L = 0$ and thus $K = I_\mu^\lambda(\mathbf{c})$ as desired. \square

4.4 Universality of the quantization

There is already a rich literature on the theory of deformation quantizations of symplectic varieties. The most relevant work for us is that of Bezrukavnikov and Kaledin [BeKa], whose show the existence and uniqueness of deformation quantizations of symplectic resolutions. This theory can be applied directly to a smooth convolution variety $\text{Gr}_\mu^{\bar{\lambda}}$. Moreover, as noted by Braden, Proudfoot and the second author [BPW, 3.4], it can be extended in a very straightforward way to the non-smooth case $\text{Gr}_\mu^{\bar{\lambda}}$, since we know that $\text{Gr}_\mu^{\bar{\lambda}}$ is a terminalization (Theorem 2.7).

This shows that the variety $\text{Gr}_\mu^{\bar{\lambda}}$ has a canonical family of quantizations which extend to a deformation quantization sheaf on $\text{Gr}_\mu^{\bar{\lambda}}$. The base of this family is the same as the base for the universal deformation of $\text{Gr}_\mu^{\bar{\lambda}}$ as a symplectic singularity (as constructed by Kaledin-Verbitsky [KV] or Namikawa [N2]). By [N3, 1.1], this base \mathbb{B} is an affine space modulo the action of a finite group. This group can be described by looking at the codimension 2 strata of the product of $\text{Gr}_\mu^{\bar{\lambda}}$, which are $\text{Gr}_\mu^{\lambda-\alpha_i}$, and taking the product of the Weyl groups attached to them by the McKay correspondence, which (using Example 2.2) in our case results in the symmetric groups $S_{\lambda,\mu} = \prod_{i: m_i > 0} S_{\lambda_i}$. Here we use the fact that these strata are simply connected.

For the remainder of this section, let us regard the complex number $r_i^{(s)}$ and $c_i^{(s)}$ as variables and let \tilde{Y}_μ be the $\mathbb{C}[r_i^{(s)}]$ -algebra which recovers the old $Y_\mu(\mathbf{r})$ upon specializing the variables. Let $\tilde{Y}_\mu^\lambda = \tilde{Y}_\mu \otimes_{\mathbb{C}[r_i^{(s)}]} \mathbb{C}[c_i^{(s)}]/(\{A_i^{(s)} : s > m_i\})$ (here we use a map $\mathbb{C}[r_i^{(s)}] \rightarrow \mathbb{C}[c_i^{(s)}]$ given by (8)). If Conjecture 2.20 (and hence Theorem 4.10) holds, then \tilde{Y}_μ^λ can be specialized (via a map $\mathbb{C}[c_i^{(s)}] \rightarrow \mathbb{C}$) to each of the $Y_\mu^\lambda(\mathbf{c})$. We conjecture that \tilde{Y}_μ^λ is related to the above universal quantization as follows.

First note that the BD analogue $\text{Gr}_{\mu; \mathbb{A}^{\rho(\lambda)}}^{\bar{\lambda}}$ is a symplectic deformation of $\text{Gr}_{\mu}^{\bar{\lambda}}$ over the base $\mathbb{A}^{\rho(\lambda)}$, and thus is the pull-back of the universal deformation by a map $b: \mathbb{A}^{\rho(\lambda)} \rightarrow \mathbb{B}$.

Conjecture 4.11. 1. The map $b: \mathbb{A}^{\rho(\lambda)} \rightarrow \mathbb{B}$ descends to a surjective map $\tilde{b}: \mathbb{A}^{\rho(\lambda)} / S_{\lambda, \mu} \rightarrow \mathbb{B}$.

2. The algebra $\tilde{Y}_{\mu}^{\lambda}$ is the base change along \tilde{b} of the universal, Bezrukavnikov-Kaledin-type quantization.

Example 4.12. We continue Example 4.6, so $G = SL_2$ and $\lambda = \alpha^{\vee}, \mu = 0$. Note that in Y_{μ}^{λ} , we have that $E^{(s)} = (-A^{(1)})^{s-1} E^{(1)}$, and $F^{(s)} = F^{(1)} (-A^{(1)})^{s-1}$, and so Y_{μ}^{λ} is generated by $E^{(1)}$ and $F^{(1)}$.

Let $U_h \mathfrak{sl}_2$ denote the h -version of the universal enveloping algebra of \mathfrak{sl}_2 . Let $C = EF + FE + \frac{1}{2}H^2$ be its Casimir element. For any complex number c , let Z_c denote the ideal in $U_h \mathfrak{sl}_2$ generated by the central element $C - c$. Standard results give that $U_h \mathfrak{sl}_2 / Z_c$ is a quantization of the nilpotent cone of \mathfrak{sl}_2 , which is isomorphic as a Poisson variety to $\text{Gr}_{\mu}^{\bar{\lambda}}$.

The map

$$E^{(1)} \mapsto E, \quad H^{(1)} \mapsto H + c^{(1)} + h, \quad F^{(1)} \mapsto F$$

defines an isomorphism $Y_{\mu}^{\lambda} \cong U_h \mathfrak{sl}_2 / Z_c$, where $c = 2c^{(2)} - \frac{1}{2}(c^{(1)})^2 + \frac{1}{2}h^2$. If we don't specialize, then the same formulas combined with the assignment

$$c^{(2)} \mapsto -\frac{1}{2}C + \frac{1}{4}(c^{(1)})^2 - \frac{1}{4}h^2$$

give an isomorphism

$$\tilde{Y}_0^2 \cong U_h(\mathfrak{sl}_2)[c^{(1)}].$$

In this example, $U_h(\mathfrak{sl}_2)$ is the universal quantization, and $c^{(1)}$ a trivial deformation parameter. The universal family is

$$\mathfrak{sl}_2 \xrightarrow{\text{tr}(a^2)} \mathbb{C}.$$

Since the fiber of the BD analogue over $(x, y) \in \mathbb{A}^2$ can be identified with matrices with eigenvalues x and y , the map b is just $b(x, y) = 1/4(x - y)^2$. Thus, choosing $x + y$ and $(x - y)^2$ as generators of symmetric functions, \tilde{b} is just the projection map $\mathbb{A}^2 \rightarrow \mathbb{A}^1$.

The sum of the $c_i^{(1)}$ is always a trivial deformation parameter; usually this is the only such parameter, but there are degenerate cases where other parameters can be trivialized as well (for example, if $\lambda = \mu$).

4.5 Quantization of Zastava spaces

In this section, we assume that Conjecture 2.20 holds and thus we will assume the conclusions of Theorem 4.10.

Let us fix ν in the positive coroot cone. Choose some μ_0 such that $\mu_0 + \nu$ is dominant. Let \mathbf{c} be a collection of complex numbers as above and consider $Y_{\mu_0}^{\mu_0 + \nu}(\mathbf{c})$.

Now for any dominant μ with $\mu \geq \mu_0$, we extend \mathbf{c} by 0 and (slightly abusing notation) consider $Y_{\mu}^{\mu + \nu}(\mathbf{c})$. Since the generators of $Y_{\mu}^{\mu + \nu}(\mathbf{c})$ are a subset of the generators of $Y_{\mu_0}^{\mu_0 + \nu}(\mathbf{c})$ and the relations are the same, we obtain a map $Y_{\mu}^{\mu + \nu}(\mathbf{c}) \rightarrow Y_{\mu_0}^{\mu_0 + \nu}(\mathbf{c})$. It is easy to see that this map is an isomorphism on the N th filtered piece if $\langle \mu, \alpha_i \rangle \geq N$ for all i .

Thus this system stabilizes to the algebra $Y_{\infty}^{\infty + \nu}$, which is the quotient of the Borel Yangian Y_{∞} by the 2-sided ideal generated by $A_i^{(s)}$ for $s > \langle \nu, \alpha_i \rangle$; perhaps surprisingly, this limit doesn't depend on \mathbf{c} or our starting μ_0 .

Combining together Theorem 4.10 with Theorem 2.10, we obtain the following (dependent on Conjecture 2.20), which was conjectured in [FR] for $G = SL_n$ (and proven for $G = SL_2$).

Theorem 4.13. $Y_{\infty}^{\infty + \nu} / \hbar Y_{\infty}^{\infty + \nu}$ is isomorphic to the Poisson algebra $\mathcal{O}(Z_{\nu})$. \square

Remark 4.14. As mentioned above, the GKLO representation gives rise to a map of graded Poisson algebras

$$Y_{\mu}^{\lambda}(\mathbf{c}) / \hbar Y_{\mu}^{\lambda}(\mathbf{c}) \rightarrow D_{\mu}^{\lambda}(\mathbf{c}) / \hbar D_{\mu}^{\lambda}(\mathbf{c})$$

(which we expect is an inclusion) and thus to a \mathbb{C}^{\times} -equivariant map of Poisson varieties

$$\prod_i (\mathbb{C}^{m_i} \setminus \Delta) \times (\mathbb{C}^{\times})^{m_i} \rightarrow \mathrm{Gr}_{\mu}^{\lambda}$$

which we expect to be étale.

If we then compose with the map $\mathrm{Gr}_{\mu}^{\lambda} \rightarrow Z_{\lambda - \mu}$, we obtain $\prod_i (\mathbb{C}^{m_i} \setminus \Delta) \rightarrow Z_{\lambda - \mu}$, which was studied in [GKLO].

References

- [Be] A. Beauville, Symplectic singularities, *Invent. Math.* **139** (2000), no. 3, 541–549.
- [BGS] A. Beilinson, V. Ginzburg, and W. Soergel, Koszul duality patterns in representation theory, *J. Amer. Math. Soc.* **9** (1996), no. 2, 473–527.
- [BeKa] R. Bezrukavnikov and D. Kaledin, Fedosov quantization in algebraic context, *Mosc. Math. J.* **4** (2004), no. 3, 559–592, 782.

- [BLPW] T. Braden, A. Licata, N. Proudfoot, and B. Webster, Quantizations of conical symplectic resolutions II: category \mathcal{O} , in preparation.
- [BPW] T. Braden, N. Proudfoot, and B. Webster, Quantizations of conical symplectic resolutions I: local and global structure; [arXiv:1208.3863](#)
- [BF] A. Braverman and M. Finkelberg, Semi-infinite Schubert varieties and quantum K-theory of flag manifolds; [arXiv:1111.2266](#).
- [BrK11] J. Brundan and A. Kleshchev, Shifted Yangians and finite W-algebras. *Adv. Math.* **200** (2006), no. 1, 136–195.
- [BrK12] ———, Representations of shifted Yangians and finite W-algebras. *Mem. Amer. Math. Soc.* **196** (2008), no. 918, viii+107 pp.
- [Dr1] V. Drinfeld, Quantum groups, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986), 798–820, Amer. Math. Soc., Providence, RI, 1987
- [Dr2] ———, On Poisson homogeneous spaces of Poisson-Lie groups. *Teoret. Mat. Fiz.* **95** (1993), no. 2, 226–227; translation in *Theoret. and Math. Phys.* **95** (1993), no. 2, 524–525.
- [EK] P. Etingof and D. Kazhdan, Quantization of Lie bialgebras III, *Selecta Math. (N.S.)* **4** (1998), no. 2, 233–269.; [arXiv:q-alg/9610030](#).
- [FKMM] M. Finkelberg, A. Kuznetsov, N. Markarian, and I. Mirković, A note on the symplectic structure on the space of G-monopoles, *Comm. Math. Phys.* **201** (1999), no. 2, 411–421.; [arXiv:math/9803124](#).
- [FM] M. Finkelberg and I. Mirković, Semiinfinite Flags. I. Case of global curve P^1 , Differential topology, infinite-dimensional Lie algebras, and applications, 81–112, Amer. Math. Soc. Transl. Ser. 2, 194, Amer. Math. Soc., Providence, RI, 1999.; [arXiv:alg-geom/9707010](#).
- [FR] M. Finkelberg and L. Rybnikov, Quantization of Drinfeld Zastava; [arXiv:1009.0676](#).
- [FZ] S. Fomin and A. Zelevinsky, Double Bruhat cells and total positivity, *JAMS*, **12**, no. 2 (1999), 335–380.
- [Ga1] F. Gavarini, The global quantum duality principle, *Journal für die reine und angewandte Mathematik*, **612** (2007), 17–33.
- [Ga2] ———, The quantum duality principle, *Annales de l’Institut Fourier*, **52**, no. 3 (2002), 809–834.

- [GKLO] A. Gerasimov, S. Kharchev, D. Lebedev, and S. Oblazin, On a class of representations of the Yangian and moduli space of monopoles; [arXiv:math/0409031](#).
- [Gi] V. Ginzburg, Perverse sheaves on a loop group and Langlands duality; [arXiv:alg-geom/9511007](#).
- [KV] D. Kaledin and M. Verbitsky, Period map for non-compact holomorphically symplectic manifolds, *Geom. Funct. Anal.* **12** (2002), no. 6, 1265–1295.
- [Lo] I. Losev, Isomorphisms of quantizations via quantization of resolutions. *Adv. Math.* **231** (2012), no. 3–4, 1216–1270
- [LY] J.-H. Lu and M. Yakimov, Group orbits and regular partitions of Poisson manifolds. *Comm. Math. Phys.* **283** (2008), no. 3, 729–748.
- [Lu] G. Lusztig, Singularities, character formulas, and a q -analog of weight multiplicities, in: *Analysis and topology on singular spaces, II, III (Luminy, 1981)*, pp. 208–229, Astérisque, vol. 101, Soc. Math. France, 1983.
- [Mir] I. Mirković, personal communication.
- [MV] I. Mirković, K. Vilonen, Geometric Langlands duality and representations of algebraic groups over commutative rings, *Ann. of Math. (2)* **166** (2007), no. 1, 95–143; [arXiv:math/0401222](#).
- [MVy] I. Mirković and M. Vybornov, Quiver varieties and Beilinson-Drinfeld Grassmannians of type A; [arXiv:0712.4160v2](#).
- [N1] Y. Namikawa, A note on symplectic singularities; [arXiv:math/0101028](#).
- [N2] ———, Poisson deformations of affine symplectic varieties. *Duke Math. J.* **156** (2011), no. 1, 5185.
- [N3] ———, Poisson deformations of affine symplectic varieties, II, *Kyoto J. Math.* **50** (2010), no. 4, 727–752.
- [R] R. Richardson, Intersections of double cosets in algebraic groups. *Indag. Math. (N.S.)* **3** (1992), no. 1, 6977.
- [W] J. Weyman, The equations of conjugacy classes of nilpotent matrices, *Invent. Math.* **98**, no. 2 (1989).